Prior Authorization Criteria Last

Updated: June 20, 2024 Effective Date: July 1, 2024



# 2024 Prior Authorizations

(List of Prior Authorizations)

PLEASE READ CAREFULLY: THIS DOCUMENT CONTAINS INFORMATION ABOUT THE PRIOR AUTHORIZATIONS ON DRUGS THAT WE COVER IN THIS PLAN.

**Note to existing members:** Beneficiaries must use network pharmacies to access their prescription drug benefit. "Benefits, List of Covered Drugs, pharmacy and provider networks and copayments may change from time to time throughout the year and on January 1 of each year."

IEHP DualChoice (HMO D-SNP) is a HMO Plan with a Medicare contract. Enrollment in IEHP DualChoice (HMO D-SNP) depends on contract renewal.

You can get this document for free in other formats, such as large print, braille, and/or audio. Call IEHP DualChoice Member Services at 1-877-273-IEHP (4347), 8am-8pm (PST), 7 days a week, including holidays. TTY users should call 1-800-718-4347. The call is free.

Puede obtener este documento gratis en otros formatos, como letra grande, Braille y/o audio. Llame a Servicios para Miembros de IEHP DualChoice al 1-877-273-IEHP (4347), 8am a 8pm (Hora del Pacífico), los 7 días de la semana, incluidos los días festivos. Los usuarios de TTY deben llamar al 1-800-718-4347. La llamada es gratuita.

您可以免費索取本文件的其他格式,例如大字版、盲文版和/或音訊版。請致電1-877-273-IEHP (4347) 與IEHP DualChoice會員服務處聯絡。服務時間為上午8點至晚上8點(太平洋標準時間),每週7天,包括節假日。TTY使用者應撥打1-800-718-4347。電話服務免費。

Quý vị có thể tải miễn phí tài liệu này ở các định dạng khác, chẳng hạn như bản in cỡ lớn, chữ nổi Braille và/hoặc tệp âm thanh. Hãy gọi Ban Dịch Vụ Hội Viên IEHP DualChoice theo số 1-877-273-IEHP (4347), 8 giờ sáng - 8 giờ tối (Múi giờ PST), 7 ngày ngày trong tuần, kể cả các ngày lễ. Người dùng TTY vui lòng gọi số 1-800-718-4347. Miễn phí cước gọi.

# **ABELCET**

### **Products Affected**

• ABELCET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Subject to Part B vs Part D determination. Failure or clinically significant adverse effects to the formulary alternative: conventional Amphotericin B.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **ABILIFY MAINTENA**

### **Products Affected**

- ABILIFY ASIMTUFII INTRAMUSCULAR SUSPENSION,EXTENDED REL SYRING 720 MG/2.4 ML, 960 MG/3.2 ML
- ABILIFY MAINTENA INTRAMUSCULAR

- SUSPENSION, EXTENDED REL RECON 300 MG, 400 MG
- ABILIFY MAINTENA INTRAMUSCULAR SUSPENSION,EXTENDED REL SYRING

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documented history of receiving oral aripiprazole without any clinically significant side effects.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to two of the formulary alternatives: Invega Sustenna, Invega Trinza or Risperdal Consta.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **ACITRETIN**

### **Products Affected**

• acitretin

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Dermatologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to two of the formulary alternatives: calcipotriene, clobetasol, cyclosporine, fluocinonide, methotrexate, or Tazorac.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **ACTIMMUNE**

### **Products Affected**

• ACTIMMUNE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Hematologist, Infectious Disease specialist, Oncologist, Orthopedist, Rheumatologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **ADEFOVIR**

### **Products Affected**

• adefovir

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Gastroenterologist, Hepatologist, Infectious Disease specialist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **ADHD**

### **Products Affected**

- dexmethylphenidate oral tablet
- methylphenidate hcl oral capsule, er biphasic 30-70
- methylphenidate hcl oral capsule,er biphasic 50-50
- methylphenidate hcl oral solution
- methylphenidate hcl oral tablet
- methylphenidate hcl oral tablet extended release

Criteria Details
Until the end of calendar year.
All FDA-approved Indications.
No

## **AIMOVIG**

### **Products Affected**

• AIMOVIG AUTOINJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Neurologist, Headache Specialist, Pain Specialist
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to one of the formulary alternatives: divalproex, valproic acid, or topiramate and failure or clinically significant adverse effects to one of the formulary alternatives: metoprolol, timolol, propranolol.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **ALOSETRON**

### **Products Affected**

alosetron

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Irritable bowel syndrome with diarrhea: Failure or clinically significant adverse effects to the formulary alternative: loperamide.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **AMBISOME**

### **Products Affected**

• AMBISOME

• amphotericin b liposome

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Subject to Part B vs Part D determination. Failure or clinically significant adverse effects to the formulary alternative: conventional Amphotericin B.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **AMITRIPTYLINE**

### **Products Affected**

• amitriptyline

- perphenazine-amitriptyline
- amitriptyline-chlordiazepoxide oral tablet

12.5-5 mg, 25-10 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	Approve if less than 65 years of age. If aged 65 years and older, prior authorization criteria will apply.
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Depression: Failure or clinically significant adverse effects to two of the formulary alternatives: citalopram, escitalopram, fluoxetine, sertraline, venlafaxine, desvenlafaxine, duloxetine, mirtazapine, or bupropion
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **AMOXAPINE**

### **Products Affected**

• amoxapine

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	Approve if less than 65 years of age. If aged 65 years and older, prior authorization criteria will apply.
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to two of the formulary alternatives: citalopram, escitalopram, fluoxetine, sertraline, venlafaxine, desvenlafaxine, duloxetine, mirtazapine, or bupropion
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **AMPHOTERICIN B**

### **Products Affected**

• amphotericin b

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Subject to Part B vs Part D determination.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **ANTICONVULSANTS 1**

### **Products Affected**

- APTIOM ORAL TABLET 200 MG, 400 MG, 600 MG, 800 MG
- FYCOMPA ORAL SUSPENSION
- FYCOMPA ORAL TABLET

• EPIDIOLEX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to one of the formulary alternatives: carbamazepine, clonazepam, divalproex, ethosuximide, felbamate, gabapentin, lamotrigine, levetiracetam, lorazepam, oxcarbazepine, phenytoin, tiagabine, topiramate, valproic acid or zonisamide.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **ANTICONVULSANTS 2**

### **Products Affected**

- clobazam oral suspension
- clobazam oral tablet
- DIACOMIT
- FINTEPLA

- MOTPOLY XR ORAL CAPSULE,EXTENDED RELEASE 24HR 100 MG, 150 MG, 200 MG
- VIMPAT ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Clobazam, Fenfluramine, or Stiripentol: Must be used as adjunctive treatment of seizure. Lacosamide: For partial seizure: Failure or clinically significant adverse effects to two of the following: carbamazepine, divalproex, felbamate, gabapentin, levetiracetam, oxcarbazepine, phenytoin, tiagabine, topiramate, valproic acid or zonisamide. For tonic-clonic seizure: Failure or clinically significant adverse effects to two of the following: lamotrigine, levetiracetam, phenytoin, or topiramate.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### ANTINEOPLASTIC

#### **Products Affected**

- abiraterone oral tablet 250 mg, 500 mg
- AFINITOR DISPERZ ORAL TABLET FOR SUSPENSION 2 MG
- AKEEGA
- ALECENSA
- ALUNBRIG
- AUGTYRO
- AYVAKIT ORAL TABLET 100 MG, 200 MG, 25 MG, 300 MG, 50 MG
- BALVERSA ORAL TABLET 3 MG, 4 MG, 5 MG
- BESREMI
- BOSULIF
- BRAFTOVI
- BRUKINSA
- CABOMETYX
- CALQUENCE (ACALABRUTINIB MAL)
- CAPRELSA
- COMETRIO
- COPIKTRA
- COTELLIC
- DAURISMO ORAL TABLET 100 MG, 25 MG
- ERIVEDGE
- ERLEADA ORAL TABLET 240 MG, 60 MG
- erlotinib oral tablet 100 mg, 150 mg, 25 mg
- everolimus (antineoplastic) oral tablet
- everolimus (antineoplastic) oral tablet for suspension 3 mg, 5 mg
- EXKIVITY
- FOTIVDA
- FRUZAQLA ORAL CAPSULE 1 MG, 5 MG
- GAVRETO
- gefitinib
- GILOTRIF
- GLEOSTINE
- IBRANCE

- ICLUSIG ORAL TABLET 10 MG, 15 MG, 30 MG, 45 MG
- IDHIFA
- imatinib oral tablet 100 mg, 400 mg
- IMBRUVICA ORAL CAPSULE 140 MG, 70 MG
- IMBRUVICA ORAL SUSPENSION
- IMBRUVICA ORAL TABLET 140 MG, 280 MG, 420 MG, 560 MG
- INLYTA ORAL TABLET 1 MG, 5 MG
- INQOVI
- INREBIC
- IRESSA
- IWILFIN
- JAKAFI
- JAYPIRCA ORAL TABLET 100 MG, 50 MG
- KISQALI FEMARA CO-PACK
- KISQALI ORAL TABLET 200 MG/DAY (200 MG X 1), 400 MG/DAY (200 MG X 2), 600 MG/DAY (200 MG X 3)
- KOSELUGO ORAL CAPSULE 10 MG, 25 MG
- KRAZATI
- lapatinib
- lenalidomide
- LENVIMA ORAL CAPSULE 10
   MG/DAY (10 MG X 1), 12 MG/DAY (4
   MG X 3), 14 MG/DAY(10 MG X 1-4 MG
   X 1), 18 MG/DAY (10 MG X 1-4 MG
   X2), 20 MG/DAY (10 MG X 2), 24
   MG/DAY(10 MG X 2-4 MG X 1), 4 MG,
   8 MG/DAY (4 MG X 2)
- LONSURF
- LORBRENA ORAL TABLET 100 MG, 25 MG
- LUMAKRAS ORAL TABLET 120 MG, 320 MG
- LYNPARZA
- LYTGOBI 12 MG DOSE (3X 4 MG TB)
- LYTGOBI 16 MG DOSE (4X 4 MG TB)
- LYTGOBI 20 MG DOSE (5X 4 MG TB)

- LYTGOBI ORAL TABLET 4 MG, 4 MG (4X 4 MG TB), 4 MG (5X 4 MG TB)
- MATULANE
- MEKINIST ORAL RECON SOLN
- MEKINIST ORAL TABLET 0.5 MG, 2 MG
- MEKTOVI
- NERLYNX
- NEXAVAR
- NINLARO
- NUBEQA
- ODOMZO
- OGSIVEO ORAL TABLET 100 MG, 150 MG, 50 MG
- OJEMDA ORAL SUSPENSION FOR RECONSTITUTION
- OJEMDA ORAL TABLET
- OJJAARA ORAL TABLET 100 MG, 150
   MG, 200 MG
- ONUREG
- ORGOVYX
- ORSERDU ORAL TABLET 345 MG, 86 MG
- PANRETIN
- pazopanib
- PEMAZYRE
- PIQRAY ORAL TABLET 200 MG/DAY (200 MG X 1), 250 MG/DAY (200 MG X1-50 MG X1), 300 MG/DAY (150 MG X 2)
- POMALYST
- PURIXAN
- QINLOCK
- RETEVMO ORAL CAPSULE 40 MG, 80 MG
- REVLIMID
- REZLIDHIA
- ROZLYTREK ORAL CAPSULE 100 MG, 200 MG
- ROZLYTREK ORAL PELLETS IN PACKET
- RUBRACA
- RYDAPT
- SCEMBLIX ORAL TABLET 20 MG, 40 MG
- sorafenib

- SPRYCEL ORAL TABLET 100 MG, 140 MG, 20 MG, 50 MG, 70 MG, 80 MG
- STIVARGA
- sunitinib malate
- SUTENT
- SYNRIBO
- TABLOID
- TABRECTA
- TAFINLAR
- TAGRISSO
- TALZENNA ORAL CAPSULE 0.1 MG, 0.25 MG, 0.35 MG, 0.5 MG, 0.75 MG, 1 MG
- TASIGNA
- TAZVERIK
- TEPMETKO
- THALOMID
- TIBSOVO
- TRUQAP
- TRUSELTIQ ORAL CAPSULE 100 MG/DAY (100 MG X 1), 125 MG/DAY (100 MG X1-25MG X1), 50 MG/DAY (25 MG X 2), 75 MG/DAY (25 MG X 3)
- TUKYSA ORAL TABLET 150 MG, 50 MG
- TURALIO ORAL CAPSULE 125 MG, 200 MG
- VANFLYTA
- VENCLEXTA
- VENCLEXTA STARTING PACK
- VERZENIO
- VITRAKVI ORAL CAPSULE 100 MG, 25 MG
- VITRAKVI ORAL SOLUTION
- VIZIMPRO
- VONJO
- VOTRIENT
- WELIREG
- XALKORI ORAL CAPSULE
- XALKORI ORAL PELLET 150 MG, 20 MG, 50 MG
- XERMELO
- XOSPATA
- XPOVIO ORAL TABLET 100 MG/WEEK (50 MG X 2), 40 MG/WEEK

(40 MG X 1), 40MG TWICE WEEK (40 MG X 2), 60 MG/WEEK (60 MG X 1), 60MG TWICE WEEK (120 MG/WEEK), 80 MG/WEEK (40 MG X 2), 80MG TWICE WEEK (160 MG/WEEK)

- XTANDI
- YONSA

- ZEJULA ORAL CAPSULE
- ZEJULA ORAL TABLET 100 MG, 200 MG, 300 MG
- ZELBORAF
- ZOLINZA
- ZYDELIG
- ZYKADIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# **APREPITANT**

### **Products Affected**

• aprepitant

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	6 months.
Other Criteria	Subject to Part B vs Part D determination. Failure or clinically significant adverse effects to one of the formulary 5-HT3 antagonist alternatives: ondansetron or granisetron except when the member is on any chemotherapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **ARCALYST**

### **Products Affected**

• ARCALYST

PA Criteria	Criteria Details
Exclusion Criteria	Concurrently taking any tumor necrosis factor (TNF)-blocking agents such as Enbrel, Humira, or Remicade.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **ARMODAFINIL**

### **Products Affected**

• armodafinil

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Narcolepsy: Failure or clinically significant adverse effects to all of the formulary alternatives: dextroamphetamine and methylphenidate.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **ATOVAQUONE**

### **Products Affected**

atovaquone

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Pneumocystic pneumonia: Failure or clinically significant adverse effects to the formulary alternative: trimethoprim/sulfamethoxazole.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **AUBAGIO**

### **Products Affected**

• AUBAGIO

• teriflunomide

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **AUSTEDO**

### **Products Affected**

• AUSTEDO ORAL TABLET 12 MG, 6 MG, 9 MG

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with an MAOI. Untreated or inadequately-treated depression, or current suicidality in patients with Huntington disease.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Neurologist, Psychiatrist
Coverage Duration	Until the end of calendar year.
Other Criteria	Chorea (Huntington's Disease): Failure or clinically significant adverse effects to the formulary alternative: tetrabenazine. Reauthorization only: Documentation of positive response to medication therapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **AUVELITY**

### **Products Affected**

• AUVELITY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Depression: Failure or clinically significant adverse effects to two of the formulary alternatives: citalopram, escitalopram, fluoxetine, sertraline, venlafaxine, desvenlafaxine, duloxetine, mirtazapine, or bupropion.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **BRONCHITOL**

### **Products Affected**

• BRONCHITOL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	New: Documentation that patient has passed the BRONCHITOL Tolerance Test (BTT). Reauthorization only: Documentation of positive response to medication therapy (improvement in lung function as determined by change in FEV1).
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **CASPOFUNGIN**

### **Products Affected**

• caspofungin

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **CIMZIA**

### **Products Affected**

• CIMZIA

### • CIMZIA POWDER FOR RECONST

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Rheumatologist, Dermatologist, or Gastroenterologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Rheumatoid Arthritis: Failure or clinically significant adverse effects to two of the formulary alternatives: Enbrel, Humira, Rinvoq, or Xeljanz IR/XR. Crohn's Disease: Failure or clinically significant adverse effects to two of the formulary alternatives: Humira, Stelara, or Skyrizi. Psoriatic arthritis: Failure or clinically significant adverse effects to two of the formulary alternatives: Enbrel, Humira, Stelara, Xeljanz IR/XR, Rinvoq, or Skyrizi. Ankylosing spondylitis: Failure or clinically significant adverse effects to two of the formulary alternatives: Enbrel, Humira, Rinvoq, or Xeljanz IR/XR. Non-radiographic Axial Spondyloarthritis: Failure or clinically significant adverse effects to the formulary alternative: Cosentyx. Plaque psoriasis: Failure or clinically significant adverse effects to two of the formulary alternatives: Humira, Stelara, Enbrel, or Skyrizi.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **CLOMIPRAMINE**

### **Products Affected**

• clomipramine

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following: Concomitant use of an MAOI, or Use within 14 days of discontinuing an MAOI, or Concomitant use of linezolid, or Concomitant use of intravenous methylene blue, or Use during the acute recovery period after a myocardial infarction.
Required Medical Information	
Age Restrictions	Approve if less than 65 years of age. If aged 65 years and older, prior authorization criteria will apply.
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Obsessive-Compulsive Disorder: Failure or clinically significant adverse effects to two of the formulary alternatives: fluoxetine, fluoxamine, or sertraline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **CONSTIPATION AGENTS**

### **Products Affected**

MOVANTIK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to one of the formulary alternatives: lactulose or polyethylene glycol.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **CORLANOR**

### **Products Affected**

- CORLANOR ORAL SOLUTION
- CORLANOR ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Heart failure in adult patients: Documented New York Association (NYHA) class II to IV heart failure with an ejection fraction of less than or equal to 35% and sinus rhythm with a resting heart rate greater than or equal to 70 beats per minute (bpm). Documentation that patient is on maximally tolerated dose of beta blocker or has a history of a documented intolerance, contraindication or a hypersensitivity to beta blocker. Documented concurrent use with an ACE inhibitor or ARB, unless both are not tolerated or contraindicated. Heart failure in pediatric patients: Documented NYHA/Ross class II to IV heart failure with an ejection fraction of less than or equal to 45% and sinus rhythm with a resting heart rate greater than or equal to 105 bpm in the age subset 6-12 months, greater than or equal to 95 bpm in the age subset 1-3 years, greater than or equal to 75 bpm in the age subset 3-5 years, greater than or equal to 70 bpm in the age subset 5-18 years
Age Restrictions	
Prescriber Restrictions	Cardiologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **COSENTYX**

### **Products Affected**

- COSENTYX (2 SYRINGES)
- COSENTYX PEN (2 PENS)
- COSENTYX SUBCUTANEOUS SYRINGE 75 MG/0.5 ML
- COSENTYX UNOREADY PEN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Plaque psoriasis: documentation of psoriasis of greater than 3% BSA or affecting crucial body areas such as hands, feet, face or genitals.
Age Restrictions	
Prescriber Restrictions	Dermatologist, Rheumatologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Ankylosing spondylitis: For adults, failure or clinically significant adverse effects to both of the formulary alternatives: Enbrel and Humira. Psoriatic arthritis: For adults, failure or clinically significant adverse effects to both of the formulary alternatives: Enbrel and Humira. Plaque Psoriasis: For adults, failure or clinically significant adverse effects to one of the formulary alternatives: Humira, Enbrel, or Skyrizi. Nonradiographic Axial Spondyloarthritis: Failure or clinically significant adverse effects to a non-steroidal anti-inflammatory drug (NSAID) or has an intolerance or contraindication to NSAID.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **CYCLOBENZAPRINE**

### **Products Affected**

• cyclobenzaprine oral tablet 10 mg, 5 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	Approve if less than 65 years of age. If aged 65 years and older, prior authorization criteria will apply.
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Documentation explaining specific benefit established with the medication, and how that benefit outweighs the potential risk
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **DALFAMPRIDINE**

### **Products Affected**

• dalfampridine

PA Criteria	Criteria Details
Exclusion Criteria	Moderate or severe renal impairment (CrCL 50 mL/min or less).
Required Medical Information	Concurrently on a disease-modifying agent for multiple sclerosis.  Documentation of difficulty walking (such as timed 25-foot walk test:  Patient must be able to walk 25 feet within 8-45 sec). Reauthorization only: Documentation of positive response to medication therapy.
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **DALIRESP**

### **Products Affected**

DALIRESP

• roflumilast

PA Criteria	Criteria Details
Exclusion Criteria	Liver impairment, moderate to severe (Child-Pugh B or C).
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to two of the formulary alternatives: either Wixela or Fluticasone/Salmeterol, Anoro Ellipta, Serevent, Spiriva or Tudorza.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **DEFERASIROX**

### **Products Affected**

• deferasirox oral tablet, dispersible

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Hematologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **DESIPRAMINE**

### **Products Affected**

• desipramine

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	Approve if less than 65 years of age. If aged 65 years and older, prior authorization criteria will apply.
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Depression: Failure or clinically significant adverse effects to two of the formulary alternatives: citalopram, escitalopram, fluoxetine, sertraline, venlafaxine, desvenlafaxine, duloxetine, mirtazapine, or bupropion
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **DIAZEPAM SOLUTION**

### **Products Affected**

• diazepam intensol

• diazepam oral solution 5 mg/5 ml (1 mg/ml)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of difficulty or inability to swallow.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **DIMETHYL FUMARATE**

### **Products Affected**

• dimethyl fumarate

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to all of the formulary alternatives: Aubagio and glatiramer.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **DISOPYRAMIDE**

### **Products Affected**

• disopyramide phosphate oral capsule

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	Approve if less than 65 years of age. If aged 65 years and older, prior authorization criteria will apply.
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Arrhythmia: Failure or clinically significant adverse effects to two of the formulary alternatives: acebutolol, flecainide, mexiletine, propafenone, quinidine, or sotalol.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **DOXEPIN**

### **Products Affected**

• doxepin oral capsule

• doxepin oral concentrate

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	Approve if less than 65 years of age. If aged 65 years and older, prior authorization criteria will apply.
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	For the average daily dose of doxepin that is greater than 6 mg: Anxiety: Failure or clinically significant adverse effects to two of the formulary alternatives: buspirone, duloxetine, escitalopram, or venlafaxine.  Depression: Failure or clinically significant adverse effects to two of the formulary alternatives: citalopram, escitalopram, fluoxetine, sertraline, venlafaxine, desvenlafaxine, duloxetine, mirtazapine, or bupropion
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **DRIZALMA**

### **Products Affected**

• DRIZALMA SPRINKLE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of difficulty or inability to swallow an intact capsule or failure or clinically significant adverse effects to the formulary alternative: duloxetine.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **DRONABINOL**

### **Products Affected**

• dronabinol

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Chemotherapy-induced nausea and vomiting: Failure or clinically significant adverse effects to two of the formulary alternatives: chlorpromazine, granisetron, metoclopramide, ondansetron, or prochlorperazine.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **DROXIDOPA**

### **Products Affected**

• droxidopa

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **DUPIXENT**

### **Products Affected**

• DUPIXENT PEN

### • DUPIXENT SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Initial: eosinophilic asthma: blood eosinophil level greater than or equal to 150 cells/mcl within the past 12 months. Eosinophilic Esophagitis: diagnosis confirmed by esophagogastroduodenoscopy with biopsy.
Age Restrictions	
Prescriber Restrictions	Initial: Atopic Dermatitis, Prurigo Nodularis: prescribed by or in consultation with a Dermatologist, Allergist or Immunologist. Asthma: prescribed by or in consultation with a physician specializing in allergy or pulmonary medicine. Chronic Rhinosinusitis with Nasal Polyposis: prescribed by or in consultation with an Otolaryngologist, Allergist or Immunologist. Eosinophilic Esophagitis: prescribed by or in consultation with a Gastroenterologist, Allergist, or Immunologist.
Coverage Duration	Until the end of calendar year.
Other Criteria	Initial: Atopic Dermatitis: 1) Atopic Dermatitis covering at least 10 percent of body surface area or Atopic Dermatitis affecting the face, head, neck, hands, feet, groin, or intertriginous areas, 2) intractable pruritus or cracking/oozing/bleeding of affected skin, 3) trial of or contraindication to one topical (corticosteroid or calcineurin inhibitor), and 4) no concurrent use with other systemic biologic/JAK inhibitor for Atopic Dermatitis.  Asthma: 1) concurrent therapy with a medium, high-dose or maximally-tolerated dose of an inhaled corticosteroid (ICS) and one other maintenance medication, 2) one asthma exacerbation requiring systemic corticosteroid burst lasting 3 or more days within the past 12 months, or one serious exacerbation requiring hospitalization or ER visit within the past 12 months, or poor symptom control despite current therapy as evidenced by at least three of the following within the past 4 weeks: daytime asthma symptoms more than twice/week, any night waking due to asthma, Short-Acting Beta Agonist (SABA) reliever for symptoms more than twice/week, any activity limitation due to asthma, and 3) no concurrent use with Xolair or other Anti-IL5 biologics when used for asthma. Chronic Rhinosinusitis with Nasal Polyposis: 1) evidence of nasal polyps by direct examination, endoscopy or sinus CT scan, 2)

PA Criteria	Criteria Details
	inadequately controlled disease as determined by use of systemic steroids in the past 2 years or endoscopic sinus surgery, and 3) a 90 day trial of one topical nasal corticosteroid. Prurigo Nodularis: 1) chronic pruritis (itch more than 6 weeks), multiple pruriginous lesions, and history or sign of a prolonged scratching behavior, 2) trial of or contraindication to one topical (corticosteroid or calcipotriene). Renewal: Atopic Dermatitis: 1) improvement while on therapy, and 2) no concurrent use with other systemic biologic/JAK inhibitor for Atopic Dermatitis. Chronic Rhinosinusitis with Nasal Polyposis, Eosinophilic Esophagitis: improvement while on therapy. Asthma: 1) no concurrent use with Xolair, or other Anti-IL5 biologics for asthma, 2) continued use of ICS and one other maintenance medication, and 3) clinical response as evidenced by: (a) reduction in asthma exacerbations from baseline, (b) decreased utilization of rescue medications, (c) increase in percent predicted FEV1 from pretreatment baseline, or (d) reduction in severity or frequency of asthma-related symptoms. Prurigo Nodularis: improvement or reduction of pruritis or pruriginous lesions.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **ELIGARD**

### **Products Affected**

- ELIGARD
- ELIGARD (3 MONTH)

- ELIGARD (4 MONTH)ELIGARD (6 MONTH)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Oncologist, Urologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **EMSAM**

### **Products Affected**

• EMSAM

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with other serotonergic drugs (i.e. SSRIs, SNRIs, TCAs)
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to all of the formulary alternatives: phenelzine and tranyleypromine.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **ENBREL**

### **Products Affected**

- ENBREL
- ENBREL MINI

### • ENBREL SURECLICK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Plaque psoriasis: documentation of psoriasis of greater than 3% BSA or affecting crucial body areas such as hands, feet, face or genitals.
Age Restrictions	
Prescriber Restrictions	Dermatologist, Rheumatologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Rheumatoid Arthritis: Failure or clinically significant adverse effects to one of the formulary alternatives: azathioprine, cyclosporine, hydroxychloroquine, leflunomide, methotrexate, or sulfasalazine.  Psoriatic arthritis: Failure or clinically significant adverse effects to the formulary alternative: methotrexate. Ankylosing spondylitis: Failure or clinically significant adverse effects to one of the formulary alternatives: celecoxib, diclofenac, indomethacin, naproxen, or sulindac. Plaque psoriasis: Failure or clinically significant adverse effects to one of the following: acitretin, cyclosporine, methotrexate or phototherapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **ENDARI**

### **Products Affected**

• ENDARI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Hematologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **ESBRIET**

### **Products Affected**

- ESBRIET ORAL CAPSULE
- MG
- ESBRIET ORAL CAPSULE

  ESBRIET ORAL TABLET 267 MG, 801

   pirfenidone oral capsule
   pirfenidone oral tablet 267 mg, 534 mg, 801 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Initial authorization: Diagnosis of idiopathic pulmonary fibrosis confirmed by the presence of usual interstitial pneumonia on high resolution computed tomography (HRCT) and/or surgical lung biopsy. Documentation of liver function tests, documentation of baseline forced vial capacity (FVC) greater than or equal to 50 percent of the predicted value AND documentation of percent predicted diffusing capacity of the lungs for carbon monoxide (%DLCO) greater than or equal to 30 percent.
Age Restrictions	
Prescriber Restrictions	Pulmonologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **ESTROGENS**

### **Products Affected**

- DUAVEE
- estradiol oral

• estradiol transdermal patch weekly

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	Approve if less than 65 years of age. If aged 65 years and older, prior authorization criteria will apply.
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Postmenopausal osteoporosis prophylaxis: Failure or clinically significant adverse effects all of the formulary alternatives: alendronic acid and risedronate. Vulvar and vaginal atrophy: Failure or clinically significant adverse effects to one of the formulary alternatives: estradiol cream or Premarin Cream.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## FENTANYL LOZENGE

### **Products Affected**

• fentanyl citrate buccal lozenge on a handle

PA Criteria	Criteria Details
Exclusion Criteria	Acute, intermittent, or postoperative pain.
Required Medical Information	Documentation of opioid tolerance taking around-the-clock opioid therapy consisting of at least 60mg of oral morphine daily, at least 25mg transdermal fentanyl/hour, at least 30 mg of oral oxycodone daily, at least 8mg oral hydromorphone daily or an equianalgesic dose of another opioid daily for a week or longer for breakthrough pain of cancer. Patients must remain on around-the clock opioids when taking transmucosal immediate release fentanyl.
Age Restrictions	
Prescriber Restrictions	Pain Specialist, Oncologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **FETZIMA**

### **Products Affected**

- FETZIMA ORAL CAPSULE,EXT REL 24HR DOSE PACK 20 MG (2)- 40 MG (26)
- FETZIMA ORAL CAPSULE,EXTENDED RELEASE 24 HR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to two of the formulary alternatives: citalopram, escitalopram, fluoxetine, sertraline, venlafaxine, desvenlafaxine, duloxetine, mirtazapine, or bupropion.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **FIRAZYR**

### **Products Affected**

• icatibant

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of hereditary angioedema (HAE), must be confirmed by blood testing.
Age Restrictions	
Prescriber Restrictions	Allergist, Immunologist, Hematologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **FIRMAGON**

### **Products Affected**

• FIRMAGON KIT W DILUENT SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Oncologist, Urologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **GENOTROPIN**

### **Products Affected**

• GENOTROPIN

### • GENOTROPIN MINIQUICK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	AGHD(initial): diagnosis confirmed as a result of past diagnosis of childhood-onset GHD, or adult-onset GHD with documentation of hormone deficiency due to hypothalamic-pituitary disease from organic or known causes (eg: damage from surgery, cranial irradiation, head trauma, subarachnoid hemorrhage) and documentation of one growth-hormone stimulant test (eg: insulin tolerance test, arginine/GHRH,glucagon,arginine) to confirm adult GHD w/corresponding peak GH values ([ITT at or below 5mcg/L],[GHRH+ARG at or below 11mcg/L if BMI less than 25kg/m2, at or below 8mcg/L if BMI at or above 25 and below 30kg/m2, or at or below 4mcg/L if BMI at or above 30kg/m2],[glucagon at or below 3mcg/L],[Arg at or below 0.4mcg/L]) or documentated deficiency of 3 anterior pituitary hormones (prolactin,ACTH,TSH,FSH/LH) and IFG-1/somatomedinC below age and gender adjusted normal range as provided by physicians lab. AGHD(reauthorization): Documentation of positive experience by the patient.
Age Restrictions	
Prescriber Restrictions	Endocrinologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **GEODON SOLUTION**

### **Products Affected**

• ziprasidone mesylate

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **GILENYA**

### **Products Affected**

• fingolimod

### • GILENYA

PA Criteria	Criteria Details
Exclusion Criteria	Class III or IV heart failure, decompensated heart failure requiring hospitalization, myocardial infarction, stroke, transient ischemic attack or unstable angina within the last 6 months. Concomitant use of Class Ia or Class III anti-arrhythmic drugs. Mobitz type II second-degree or third-degree atrioventricular block, or sick-sinus syndrome unless the patient has a functional pacemaker. QT interval at baseline 500 ms or greater.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	Until the end of calendar year.
Other Criteria	For adults: Failure or clinically significant adverse effects to Aubagio and glatiramer.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **GLATIRAMER**

### **Products Affected**

• glatiramer

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **GLATOPA**

### **Products Affected**

• glatopa

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### GLP1

#### **Products Affected**

- OZEMPIC SUBCUTANEOUS PEN INJECTOR 0.25 MG OR 0.5 MG (2 MG/3 ML), 0.25 MG OR 0.5 MG(2 MG/1.5 ML), 1 MG/DOSE (4 MG/3 ML), 2 MG/DOSE (8 MG/3 ML)
- TRULICITY
- VICTOZA SUBCUTANEOUS PEN INJECTOR 0.6 MG/0.1 ML (18 MG/3 ML)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **GLYBURIDE**

### **Products Affected**

- glyburide
- glyburide micronized

• glyburide-metformin oral tablet 1.25-250 mg, 2.5-500 mg, 5-500 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	Approve if less than 65 years of age. If aged 65 years and older, prior authorization criteria will apply.
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to formulary alternative: glipizide.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **GUANFACINE**

### **Products Affected**

• guanfacine oral tablet

• guanfacine oral tablet extended release 24 hr

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	Approve if less than 65 years of age. If aged 65 years and older, prior authorization criteria will apply.
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Hypertension: Failure or clinically significant adverse effects to two of the formulary alternatives: benazepril, fosinopril, hydrochlorothiazide, irbesartan, lisinopril, losartan, losartan/hydrochlorothiazide, lisinopril/hydrochlorothiazide, quinapril/hydrochlorothiazide, quinapril, ramipril, or valsartan/hydrochlorothiazide. ADHD: Failure or clinically significant adverse effects to two of the formulary alternatives: amphetamine/dextroamphetamine, dexmethylphenidate, dextroamphetamine, or methylphenidate.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **HP ACTHAR**

### **Products Affected**

• ACTHAR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Neurologist for infantile spasm and exacerbation of multiple sclerosis
Coverage Duration	Multiple sclerosis: 21 days. For other approved indications: 28 days.
Other Criteria	For acute exacerbations of multiple sclerosis, patients must be receiving concurrent immunomodulator therapy, such as Aubagio, glatiramer, or interferon beta 1a. For all other non-neurological indications, failure or clinically significant adverse effects to other first line or standard of care therapies must be submitted.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **HRM ANTIPSYCHOTICS**

### **Products Affected**

• molindone oral tablet 10 mg, 25 mg, 5 mg • thioridazine

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	Approve if less than 65 years of age. If aged 65 years and older, prior authorization criteria will apply.
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to two of the formulary alternatives: olanzapine, risperidone, quetiapine, ziprasidone, or aripiprazole.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **HUMIRA**

#### **Products Affected**

- HUMIRA PEN
- HUMIRA PEN CROHNS-UC-HS START
- HUMIRA SUBCUTANEOUS SYRINGE KIT 40 MG/0.8 ML
- HUMIRA(CF)
- HUMIRA(CF) PEN
- HUMIRA(CF) PEN CROHNS-UC-HS
- HUMIRA(CF) PEN PSOR-UV-ADOL HS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Plaque psoriasis: documentation of psoriasis involving 3% of BSA or greater, or affecting crucial body areas such as hands, feet, face or genitals.
Age Restrictions	
Prescriber Restrictions	Dermatologist, Gastroenterologist, Ophthalmologist, Rheumatologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Rheumatoid Arthritis: Failure or clinically significant adverse effects to one of the formulary alternatives: azathioprine, cyclosporine, hydroxychloroquine, leflunomide, methotrexate, or sulfasalazine.  Psoriatic arthritis: Failure or clinically significant adverse effects to the formulary alternative: methotrexate. Ankylosing spondylitis: Failure or clinically significant adverse effects to one of the formulary alternatives: celecoxib, diclofenac, indomethacin, naproxen, or sulindac. Plaque psoriasis: Failure or clinically significant adverse effects to one of the formulary alternatives: acitretin, cyclosporine, methotrexate or phototherapy. Crohn's disease and Ulcerative colitis: Failure or clinically significant adverse effects to one of the formulary alternatives: budesonide, mesalamine or sulfasalazine.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **HUMIRA PEDIATRIC**

#### **Products Affected**

- HUMIRA(CF) PEDI CROHNS STARTER
- HUMIRA(CF) PEN PEDIATRIC UC

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **HUMIRA PSORIASIS**

### **Products Affected**

• HUMIRA PEN PSOR-UVEITS-ADOL HS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Plaque psoriasis: documentation of psoriasis involving 3% of BSA or greater, or affecting crucial body areas such as hands, feet, face or genitals.
Age Restrictions	
Prescriber Restrictions	Dermatologist, Rheumatologist, Ophthalmologist, Gastroenterologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **IMIPRAMINE**

### **Products Affected**

• imipramine hcl

• imipramine pamoate

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	Approve if less than 65 years of age. If aged 65 years and older, prior authorization criteria will apply.
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **INCRELEX**

### **Products Affected**

• INCRELEX

PA Criteria	Criteria Details
Exclusion Criteria	Pediatric patients with malignant neoplasm or history of malignancy. Use for growth promotion in patients with closed epiphyses.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Endocrinologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **INDOMETHACIN**

### **Products Affected**

• indomethacin oral capsule

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	Approve if less than 65 years of age. If aged 65 years and older, prior authorization criteria will apply.
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to two of the formulary alternatives: celecoxib, diclofenac, ibuprofen, meloxicam, nabumetone, naproxen, or sulindac.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **INGREZZA**

### **Products Affected**

• INGREZZA

- INGREZZA SPRINKLE
- INGREZZA INITIATION PK(TARDIV)

INUKEZZA INITIATION PK(TARDIV)	
PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Failure or clinically significant adverse effects to the formulary alternative: Austedo. Documentation of baseline Abnormal Involuntary Movement Scale (AIMS) scores. Reauthorization only: Documentation of positive response to medication therapy as evidenced by an improved AIMS score as compared to baseline.
Age Restrictions	
Prescriber Restrictions	Neurologist, Psychiatrist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **INTERFERON BETA-1A**

- PLEGRIDY SUBCUTANEOUS PEN INJECTOR 125 MCG/0.5 ML
- PLEGRIDY SUBCUTANEOUS SYRINGE 125 MCG/0.5 ML
- REBIF (WITH ALBUMIN)
- REBIF REBIDOSE
- REBIF TITRATION PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to all of the formulary alternatives: Aubagio and glatiramer.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **INVEGA HAFYERA**

### **Products Affected**

• INVEGA HAFYERA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to all of the formulary alternatives: oral paliperidone and Invega Sustenna.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **INVEGA SUSTENNA**

### **Products Affected**

• INVEGA SUSTENNA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to the formulary alternative: oral paliperidone.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **INVEGA TRINZA**

### **Products Affected**

• INVEGA TRINZA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to all of the formulary alternatives: oral paliperidone and Invega Sustenna.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **ITRACONAZOLE**

### **Products Affected**

• itraconazole oral solution

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **JYLAMVO**

### **Products Affected**

• JYLAMVO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Oncologist, Dermatologist, Hematologist , Rheumatologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Subject to Part B vs Part D determination.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **KALYDECO**

### **Products Affected**

• KALYDECO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of a mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to ivacaftor.
Age Restrictions	
Prescriber Restrictions	Infectious Disease specialist, Pulmonologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **KERENDIA**

### **Products Affected**

• KERENDIA

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with a strong CYP3A4 inhibitor or diagnosis of adrenal insufficiency
Required Medical Information	Labs within the past 30 days documenting serum potassium level of less than or equal to 5.0 mEq/L, estimated glomerular filtration rate of at least 25 mL/min/1.73m2 and urine albumin-to-creatinine ratio (UACR) of at least 30 mg/g (2) Receiving concurrent therapy with angiotensin-converting enzyme inhibitor (ACE inhibitor) or angiotensin receptor blocker (ARB) at maximally tolerated labeled dosage, unless contraindicated (3) medical justification that a sodium-glucose cotransport-2 (SGLT2) inhibitor (Jardiance, Invokana, Farxiga, Steglatro) AND a steroidal mineralocorticoid receptor antagonist (spironolactone, eplerenone) have been tried and failed, are contraindicated, or would not be medically appropriate for the patient.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **KINERET**

### **Products Affected**

• KINERET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Rheumatoid Arthritis: Failure or clinically significant adverse effects to two of the formulary alternatives: Enbrel, Humira, Rinvoq, or Xeljanz IR/XR.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# **KORLYM**

### **Products Affected**

• KORLYM

• mifepristone oral tablet 300 mg

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use of simvastatin, lovastatin and CYP3A substrates with narrow therapeutic ranges (e.g. cyclosporine, fentanyl, sirolimus, etc.). History of unexplained vaginal bleeding or endometrial hyperplasia with atypia or endometrial carcinoma.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### LEDIPASVIR-SOFOSBUVIR

### **Products Affected**

• ledipasvir-sofosbuvir

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of chronic hepatitis C infection confirmed by a detectable serum hepatitis C virus RNA through quantitative assay. Documentation of genotype. Documentation of the absence or presence of cirrhosis and if compensated or decompensated. Documentation of any previous treatment. Documentation of liver transplant status. Treatment regimen will be approved based on genotype and previous treatment experience as defined by current AASLD guidelines.
Age Restrictions	
Prescriber Restrictions	Gastroenterologist, Hepatologist, Infectious Disease specialist
Coverage Duration	Duration will be applied consistent with current AASLD/IDSA guidance.
Other Criteria	Hepatitis C: Failure or clinically significant adverse effects to the formulary alternative: sofosbuvir-velpatasvir (generic Epclusa).
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **LEUKINE**

### **Products Affected**

• LEUKINE INJECTION RECON SOLN

PA Criteria	Criteria Details
Exclusion Criteria	Excessive leukemia myeloid blasts in the bone marrow or peripheral blood equal to or greater than 10%.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Hematologist, Oncologist
Coverage Duration	3 months.
Other Criteria	Failure or clinically significant adverse effects to formulary alternative: Nivestym and Zarxio.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## LEUPROLIDE ACETATE

### **Products Affected**

• leuprolide (3 month)

• leuprolide subcutaneous kit

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Oncologist, Urologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Subject to Part B vs Part D determination.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **LEVALBUTEROL**

#### **Products Affected**

• levalbuterol hcl inhalation solution for nebulization 0.31 mg/3 ml, 0.63 mg/3 ml, 1.25 mg/0.5 ml, 1.25 mg/3 ml

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Subject to Part B vs Part D determination. Failure or clinically significant adverse effects to the formulary alternative: albuterol inhalant solution.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## LIDOCAINE PATCH

### **Products Affected**

• lidocaine topical adhesive patch,medicated 5 %

• tridacaine

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of one of the following: Pain associated with diabetic neuropathy, Pain associated with cancer-related neuropathy, Post-herpetic neuralgia, Chronic back pain, or Osteoarthritis of the knee or hip.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

### **LUPRON DEPOT**

- LUPRON DEPOT
- LUPRON DEPOT (3 MONTH)
- LUPRON DEPOT (4 MONTH)LUPRON DEPOT (6 MONTH)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Endometriosis: Patient has had surgical ablation to prevent recurrence, or history of failure, contraindication, or intolerance to oral contraceptives.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Subject to Part B vs Part D determination.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **LYRICA**

- pregabalin oral capsule 100 mg, 150 mg, 200 mg, 225 mg, 25 mg, 300 mg, 50 mg, 75 mg
  - pregabalin oral solution

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Postherpetic neuralgia: Failure or clinically significant adverse effects to the formulary alternative: gabapentin. Diabetic neuropathy: Failure or clinically significant adverse effects to all of the formulary alternatives: duloxetine and gabapentin. Fibromyalgia: Failure or clinically significant adverse effects to two of the formulary alternatives: duloxetine, gabapentin or Savella.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **MEGESTROL**

- megestrol oral suspension 400 mg/10 ml
   megestrol oral tablet (40 mg/ml)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	Approve if less than 65 years of age. If aged 65 years and older, prior authorization criteria will apply.
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Cachexia associated with AIDS: Failure or clinically significant adverse effects to all of the formulary alternatives: dronabinol and oxandrolone.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

### **MEPROBAMATE**

### **Products Affected**

• meprobamate

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	Approve if less than 65 years of age. If aged 65 years and older, prior authorization criteria will apply.
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to at least two of the formulary alternatives: buspirone, duloxetine, escitalopram, or venlafaxine.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **METHOCARBAMOL**

### **Products Affected**

• methocarbamol oral

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	Approve if less than 65 years of age. If aged 65 years and older, prior authorization criteria will apply.
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Documentation explaining specific benefit established with the medication, and how that benefit outweighs the potential risk
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **METHOXSALEN**

### **Products Affected**

methoxsalen

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Dermatologist, Rheumatologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to two of the formulary alternatives: calcipotriene, clobetasol, cyclosporine, fluocinonide, methotrexate, or tazarotene.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## METHYLDOPA/HYDROCHLOROTHIAZIDE

### **Products Affected**

• methyldopa-hydrochlorothiazide

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	Approve if less than 65 years of age. If aged 65 years and older, prior authorization criteria will apply.
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to two of the formulary alternatives: benazepril, fosinopril, hydrochlorothiazide, irbesartan, lisinopril, losartan, losartan/hydrochlorothiazide, lisinopril/hydrochlorothiazide, quinapril/hydrochlorothiazide, quinapril, ramipril, or valsartan/hydrochlorothiazide.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **METYROSINE**

### **Products Affected**

• metyrosine

PA Criteria	Criteria Details
Exclusion Criteria	Essential hypertension.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **MODAFINIL**

### **Products Affected**

• modafinil

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Narcolepsy: Failure or clinically significant adverse effects to all of the formulary alternatives: dextroamphetamine and methylphenidate.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **NAYZILAM**

### **Products Affected**

• LIBERVANT

### • NAYZILAM

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **NIVESTYM**

### **Products Affected**

• NIVESTYM INJECTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Hematologist, Infectious Disease specialist, Oncologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **NUCALA**

### **Products Affected**

• NUCALA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Allergist, Immunologist, Pulmonologist, Rheumatologist, Hematologist, or Otolaryngologist.
Coverage Duration	Until the end of calendar year.
Other Criteria	Asthma: Failure or clinically significant adverse effects to two of the formulary alternatives: 1) budesonide, Flovent, Arnuity Ellipta or Qvar and 2) fluticasone-salmeterol, Wixela, or Breo Ellipta.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **NUEDEXTA**

### **Products Affected**

• NUEDEXTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **NUPLAZID**

### **Products Affected**

NUPLAZID

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **OCTREOTIDE**

### **Products Affected**

• octreotide acetate injection solution

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **OFEV**

### **Products Affected**

• OFEV

PA Criteria	Criteria Details
Exclusion Criteria	Moderate or severe hepatic impairment (Child-Pugh B or C).
Required Medical Information	Initial authorization: Documentation of liver function tests.  Reauthorization only: Documentation of positive response to therapy and documentation of liver function tests.
Age Restrictions	
Prescriber Restrictions	Pulmonologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **OLANZAPINE ODT**

### **Products Affected**

• olanzapine oral tablet, disintegrating

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of difficulty or inability to swallow or failure or clinically significant adverse effects to the formulary alternative: oral olanzapine.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **OLANZAPINE SOLUTION**

### **Products Affected**

• olanzapine intramuscular

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **OMNITROPE**

### **Products Affected**

• OMNITROPE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	AGHD (initial): diagnosis confirmed as a result of past diagnosis of childhood-onset GHD, or adult-onset GHD with documentation of hormone deficiency due to hypothalamic-pituitary disease from organic or known causes (eg: damage from surgery, cranial irradiation, head trauma, subarachnoid hemorrhage) and documentation of one growth-hormone stimulant test (eg: insulin tolerance test, arginine/GHRH, glucagon, arginine) to confirm adult GHD w/corresponding peak GH values ([ITT at or below 5mcg/L],[GHRH+ARG at or below 11mcg/L if BMI less than 25kg/m2, at or below 8mcg/L if BMI at or above 25 and below 30kg/m2, or at or below 4mcg/L if BMI at or above 30kg/m2],[glucagon at or below 3mcg/L],[Arg at or below 0.4mcg/L]) or documented deficiency of 3 anterior pituitary hormones (prolactin, ACTH, TSH, FSH/LH) and IFG-1/somatomedin C below age and gender adjusted normal range as provided by physicians lab. AGHD (reauthorization): Documentation of positive experience by the patient.
Age Restrictions	
Prescriber Restrictions	Endocrinologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **ORAL ANTIPSYCHOTICS**

- asenapine maleate
- CAPLYTA
- FANAPT
- LATUDA ORAL TABLET 120 MG, 20 MG, 40 MG, 60 MG, 80 MG
- lurasidone oral tablet 120 mg, 20 mg, 40 mg, 60 mg, 80 mg
- VRAYLAR ORAL CAPSULE
- VRAYLAR ORAL CAPSULE,DOSE PACK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to two of the formulary alternatives: olanzapine, quetiapine, risperidone, ziprasidone or aripiprazole.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **ORENCIA**

#### **Products Affected**

• ORENCIA CLICKJECT

• ORENCIA SUBCUTANEOUS SYRINGE 125 MG/ML, 50 MG/0.4 ML, 87.5 MG/0.7 ML

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Rheumatologist, transplant specialist
Coverage Duration	Until the end of calendar year.
Other Criteria	Rheumatoid Arthritis: Failure or clinically significant adverse effects to one of the formulary alternatives: azathioprine, cyclosporine, hydroxychloroquine, leflunomide, methotrexate, or sulfasalazine. Psoriatic arthritis: Failure or clinically significant adverse effects to the formulary alternative: methotrexate.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **ORKAMBI**

#### **Products Affected**

• ORKAMBI

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of homozygous F508del mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene.
Age Restrictions	
Prescriber Restrictions	Infectious Disease specialist, Pulmonologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **OTEZLA**

#### **Products Affected**

OTEZLA

• OTEZLA STARTER ORAL TABLETS,DOSE PACK 10 MG (4)-20 MG (4)-30 MG (47)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Plaque psoriasis: documentation of psoriasis with BSA of 2% or greater or affecting crucial body areas such as hands, feet, face or genitals.
Age Restrictions	
Prescriber Restrictions	Dermatologist, Rheumatologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Psoriatic arthritis: Failure or clinically significant adverse effects to the formulary alternative: methotrexate. Plaque psoriasis: Failure or clinically significant adverse effects to one of the formulary alternatives: acitretin, cyclosporine, methotrexate or phototherapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **OXANDROLONE**

#### **Products Affected**

• oxandrolone

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **OXBRYTA**

#### **Products Affected**

- OXBRYTA ORAL TABLET 300 MG, 500 MG
- OXBRYTA ORAL TABLET FOR SUSPENSION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of at least 1 episode of vaso-occlusive crisis (VOC) in the past 12 months. Hemoglobin (Hgb) greater than or equal to 5.5 and less than or equal to 10.5 g/dL. Re-authorization: Documentation of positive response to therapy.
Age Restrictions	
Prescriber Restrictions	Hematologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **OXERVATE**

#### **Products Affected**

• OXERVATE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of Stage 2 or 3 neurotrophic keratitis.
Age Restrictions	
Prescriber Restrictions	Ophthalmologist
Coverage Duration	8 weeks.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **OXTELLAR**

#### **Products Affected**

• OXTELLAR XR ORAL TABLET EXTENDED RELEASE 24 HR 150 MG, 300 MG, 600 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to oxcarbazepine immediate release.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **PAH**

#### **Products Affected**

• ambrisentan

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Cardiologist, Pulmonologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to the formulary alternative: sildenafil or tadalafil. Approve if in combination with tadalafil for treatment naive PAH patients with WHO FC II and III.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## PARATHYROID HORMONE ANALOGS

#### **Products Affected**

• teriparatide subcutaneous pen injector 20 • TYMLOS mcg/dose (620mcg/2.48ml)

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Bone mineral density (BMD) T score of -3.5 or less based on BMD measurements from lumbar spine (at least two vertebral bodies), hip (femoral neck, total hip), or radius (one-third radius site) OR BMD T-score between -2.5 and -3.5 (BMD T-score greater than -3.5 and less than or equal to -2.5) based on BMD measurements from lumbar spine (at least two vertebral bodies), hip (femoral neck, total hip), or radius (one-third radius site) and a history of fractures.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to one of the formulary alternatives: alendronic acid or risedronate. Medical justification required for treatment duration beyond 24 months.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **PDE5 INHIBITORS**

#### **Products Affected**

- alyq
- alyqtadalafil (pulm. hypertension)TADLIQ

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with nitrates or PDE5 inhibitors.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Cardiologist, Pulmonologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **PEGFILGRASTIM**

#### **Products Affected**

- FULPHILA
- UDENYCA
- UDENYCA AUTOINJECTOR
- UDENYCA ONBODY
- **ZIEXTENZO**

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Hematologist, Infectious Disease specialist, Oncologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to formulary alternative: Nivestym and Zarxio.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **PHENOBARBITAL**

#### **Products Affected**

• phenobarbital

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	Approve if less than 65 years of age. If aged 65 years and older, prior authorization criteria will apply.
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Anticonvulsant: Failure or clinically significant adverse effects to two of the formulary alternatives: carbamazepine, divalproex, ethosuximide, felbamate, gabapentin, lamotrigine, levetiracetam, oxcarbazepine, phenytoin, tiagabine, topiramate, or zonisamide.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **PIMECROLIMUS**

#### **Products Affected**

• pimecrolimus

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to one of the topical formulary alternatives: clobetasol, betamethasone, fluocinolone or fluocinonide and failure or clinically significant adverse effects to the formulary alternative: tacrolimus ointment.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **POSACONAZOLE**

#### **Products Affected**

- NOXAFIL ORAL SUSPENSION
- posaconazole oral

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to one of the formulary alternatives: fluconazole, itraconazole, or voriconazole.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## PRADAXA PELLET

#### **Products Affected**

• PRADAXA ORAL PELLETS IN PACKET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	Pediatric patients aged 3 months to less than 12 years of age.
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## PREMARIN TABLETS

#### **Products Affected**

• PREMARIN ORAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Postmenopausal osteoporosis prophylaxis: Failure or clinically significant adverse effects to all of the formulary alternatives: alendronic acid and risedronate. Vulvar and vaginal atrophy: Failure or clinically significant adverse effects to the formulary alternative: estradiol cream. Other indication(s): Failure or clinically significant adverse effects to one of the formulary alternatives: estradiol transdermal patch or estradiol tablet.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## PREMPRO TABLETS

#### **Products Affected**

• PREMPRO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Postmenopausal osteoporosis prophylaxis: Failure or clinically significant adverse effects to all of the formulary alternatives: alendronic acid and risedronate. Vulvar and vaginal atrophy: Failure or clinically significant adverse effects to the formulary alternative: estradiol cream. Other indication(s): Failure or clinically significant adverse effects to one of the formulary alternatives: estradiol transdermal patch or estradiol tablet.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **PREVYMIS**

#### **Products Affected**

• PREVYMIS ORAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Infectious Disease specialist, Oncologist, Transplant specialist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **PROCRIT**

#### **Products Affected**

• PROCRIT INJECTION SOLUTION 10,000 UNIT/ML, 2,000 UNIT/ML,

20,000 UNIT/ML, 3,000 UNIT/ML, 4,000 UNIT/ML, 40,000 UNIT/ML

PA Criteria	Criteria Details
Exclusion Criteria	Uncontrolled hypertension.
Required Medical Information	For anemia due to chronic kidney disease: Hemoglobin (Hgb) is less than 10g/dL and documentation of transferrin saturation greater than or equal to 20% and ferritin greater than or equal to 100ng/mL. For anemia due to chemotherapy: Hemoglobin (Hgb) is less than 10g/dL. For surgical FDA indications: Hemoglobin (Hgb) is 10g/dL-13g/dL and patient is not a candidate for autologous blood donation and significant blood loss is anticipated from elective, non cardiac, or nonvascular surgery. Zidovudine induced: Hemoglobin (Hgb) is less than 11g/dL. Myelodysplastic syndrome: Hemoglobin (Hgb) is less than 11g/dL and erythropoietin is less than or equal to 500 mU/mL.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Subject to Part B vs Part D determination.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

## PROLASTIN C

#### **Products Affected**

• PROLASTIN-C INTRAVENOUS SOLUTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Clinical evidence of emphysema
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year
Other Criteria	Pre-treatment serum levels of alpha-1 antitrypsin (AAT) that is less than 11 micromol/L (80 mg/dl by radial immunodiffusion or 50 mg/dl by nephelometry)
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **PROLIA**

#### **Products Affected**

• PROLIA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of bone mineral density (BMD) T-score of -3.5 or less based on BMD measurements from lumbar spine (at least two vertebral bodies), hip (femoral neck, total hip), or radius (one-third radius site) OR BMD T-score between -2.5 and -3.5 (greater than -3.5 and less than or equal to -2.5) based on BMD measurements from lumbar spine (at least two vertebral bodies), hip (femoral neck, total hip), or radius (one-third radius site) and a history of fractures.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to all of the formulary alternatives: alendronic acid and risedronate.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **PROMACTA**

#### **Products Affected**

• PROMACTA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of platelet count. Thrombocytopenia in hepatitis C infection: Documentation of concurrent or planned interferon-based treatment of chronic hepatitis C.
Age Restrictions	
Prescriber Restrictions	Gastroenterologist, Hematologist, Hepatologist, Infectious Disease specialist, Oncologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Chronic immune (idiopathic) thrombocytopenia: Failure or clinically significant adverse effects to one of the formulary alternatives: dexamethasone, methylprednisolone, prednisolone or prednisone.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **PROMETHAZINE**

#### **Products Affected**

- promethazine oral
- promethazine rectal suppository 12.5 mg, 25 mg
- promethegan rectal suppository 25 mg, 50 mg

23 mg	
PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	Approve if less than 65 years of age. If aged 65 years and older, prior authorization criteria will apply.
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Allergy: Failure or clinically significant adverse effects to one of the formulary alternatives: cetirizine and levocetirizine. Nausea and vomiting: Failure or clinically significant adverse effects to two of the formulary alternatives: chlorpromazine, granisetron, ondansetron, or prochlorperazine.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **PROTRIPTYLINE**

#### **Products Affected**

• protriptyline

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	Approve if less than 65 years of age. If aged 65 years and older, prior authorization criteria will apply.
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to two of the formulary alternatives: citalopram, escitalopram, fluoxetine, sertraline, venlafaxine, desvenlafaxine, duloxetine, mirtazapine, or bupropion
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **PYRIMETHAMINE**

#### **Products Affected**

• pyrimethamine

PA Criteria	Criteria Details
Exclusion Criteria	Anemia due to folate deficiency
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Hematologist, HIV specialist, Infectious Disease specialist, Oncologist, Transplant specialist
Coverage Duration	Until the end of calendar year.
Other Criteria	Primary prophylaxis of toxoplasmic encephalitis: Failure or clinically significant adverse effects to the formulary alternative: trimethoprim/sulfamethoxazole.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# **QUININE**

#### **Products Affected**

• quinine sulfate

PA Criteria	Criteria Details
Exclusion Criteria	Prevention or treatment of nocturnal leg cramps.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	10 days.
Other Criteria	Failure or clinically significant adverse effects to one of the formulary alternatives: chloroquine or hydroxychloroquine.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **REPATHA**

#### **Products Affected**

- REPATHA PUSHTRONEX
- REPATHA SURECLICK

• REPATHA SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Primary hyperlipidemia: Documentation of current LDL levels above 100mg/dL while taking maximally tolerated statin therapy and ezetimibe therapy, unless intolerant or contraindicated to statin or ezetimibe therapy. Secondary prevention of ASCVD: Documentation of at least one high risk feature: recent ACS (within the past 12 months), history of MI, history of ischemic stroke, or symptomatic peripheral arterial disease (history of claudication with ABI greater than 0.85, or previous revascularization or amputation). Heterozygous Familial Hypercholesterolemia (HeFH) or Homozygous Familial Hypercholesterolemia (HoFH): Documentation to confirm diagnosis by genetic testing or by clinical criteria (such as Simon Broome or the Dutch Lipid Clinic Network criteria, or history of untreated LDL-C greater than 180 mg/dL together with xanthoma or cornealis), or evidence of Familial Hypercholesterolemia in first or second-degree relatives.
Age Restrictions	
Prescriber Restrictions	Cardiologist, Endocrinologist, Lipid specialist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **RETACRIT**

#### **Products Affected**

• RETACRIT INJECTION SOLUTION 10,000 UNIT/ML, 2,000 UNIT/ML,

20,000 UNIT/ML, 3,000 UNIT/ML, 4,000 UNIT/ML, 40,000 UNIT/ML

PA Criteria	Criteria Details
Exclusion Criteria	Uncontrolled hypertension.
Required Medical Information	For anemia due to chronic kidney disease: Hemoglobin (Hgb) is less than 10g/dL and documentation of transferrin saturation greater than or equal to 20% and ferritin greater than or equal to 100ng/mL. For anemia due to chemotherapy: Hemoglobin (Hgb) is less than 10g/dL. For surgical FDA indications: Hemoglobin (Hgb) is 10g/dL-13g/dL and patient is not a candidate for autologous blood donation and significant blood loss is anticipated from elective, non cardiac, or nonvascular surgery. Zidovudine induced: Hemoglobin (Hgb) is less than 11g/dL. Myelodysplastic syndrome: Hemoglobin (Hgb) is less than 11g/dL and erythropoietin is less than or equal to 500 mU/mL.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Subject to Part B vs Part D determination.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

### **REXULTI**

#### **Products Affected**

• REXULTI ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Schizophrenia: Failure or clinically significant adverse effects to two of the formulary alternatives: olanzapine, risperidone, quetiapine, ziprasidone or aripiprazole. Depression: Failure or clinically significant adverse effects to two of the formulary alternatives: citalopram, escitalopram, fluoxetine, mirtazapine, paroxetine, sertraline, or venlafaxine.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **REZUROCK**

#### **Products Affected**

• REZUROCK

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **RINVOQ**

#### **Products Affected**

• RINVOQ ORAL TABLET EXTENDED RELEASE 24 HR 15 MG, 30 MG, 45 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Rheumatologist, dermatologist, gastroenterologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Rheumatoid Arthritis: Failure or clinically significant adverse effects to one of the formulary alternatives: azathioprine, cyclosporine, hydroxychloroquine, leflunomide, methotrexate, or sulfasalazine. Atopic dermatitis: Failure or clinically significant adverse effects to two of the formulary alternatives: cyclosporine, azathioprine, methotrexate.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### RISPERDAL CONSTA

#### **Products Affected**

• RISPERDAL CONSTA

• UZEDY

• risperidone microspheres

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to the formulary alternative: oral risperidone.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **RUFINAMIDE**

#### **Products Affected**

- rufinamide oral suspension
- rufinamide oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Failure or clinically significant adverse effects to one of the formulary alternatives: carbamazepine, clonazepam, divalproex, ethosuximide, felbamate, gabapentin, lamotrigine, levetiracetam, lorazepam, oxcarbazepine, phenytoin, tiagabine, topiramate, clobazam or zonisamide. For Rufinamide suspension: Failure or clinically significant adverse effects to one of the formulary alternatives: Rufinamide tablet.
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **SAPROPTERIN**

#### **Products Affected**

• sapropterin

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **SECUADO**

#### **Products Affected**

• SECUADO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to two of the formulary alternatives: olanzapine, quetiapine, risperidone, ziprasidone or aripiprazole.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **SIGNIFOR**

#### **Products Affected**

• SIGNIFOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **SIRTURO**

### **Products Affected**

• SIRTURO ORAL TABLET 100 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Infectious Disease specialist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **SKYRIZI**

#### **Products Affected**

- SKYRIZI SUBCUTANEOUS PEN INJECTOR
- SKYRIZI SUBCUTANEOUS SYRINGE 150 MG/ML

• SKYRIZI SUBCUTANEOUS WEARABLE INJECTOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Plaque psoriasis: documentation of psoriasis of greater than 3% BSA or affecting crucial body areas such as hands, feet, face or genitals.
Age Restrictions	
Prescriber Restrictions	Dermatologist, gastroenterologist, rheumatologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Plaque psoriasis: Failure or clinically significant adverse effects to one of the following: acitretin, cyclosporine, methotrexate or phototherapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# SOFOSBUVIR-VELPATASVIR

### **Products Affected**

• sofosbuvir-velpatasvir

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of chronic hepatitis C infection confirmed by a detectable serum hepatitis C virus RNA through quantitative assay. Documentation of genotype. Documentation of the absence or presence of cirrhosis and if compensated or decompensated. Documentation of any previous treatment. Documentation of liver transplant status. Treatment regimen will be approved based on genotype and previous treatment experience as defined by current AASLD guidelines.
Age Restrictions	
Prescriber Restrictions	Gastroenterologist, Hepatologist, Infectious Disease specialist
Coverage Duration	Duration will be applied consistent with current AASLD/IDSA guidance.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **SOMAVERT**

### **Products Affected**

• SOMAVERT

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Endocrinologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to the formulary alternative: octreotide.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **SPRITAM**

### **Products Affected**

• SPRITAM

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to the formulary alternative: levetiracetam oral solution.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **STELARA**

### **Products Affected**

• STELARA SUBCUTANEOUS

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Plaque psoriasis: documentation of psoriasis of greater than 3% BSA or affecting crucial body areas such as hands, feet, face or genitals.
Age Restrictions	
Prescriber Restrictions	Dermatologist, Gastroenterologist, Rheumatologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Psoriatic arthritis: Failure or clinically significant adverse effects to the formulary alternative: methotrexate. Plaque psoriasis: Failure or clinically significant adverse effects to one of the formulary alternatives: acitretin, cyclosporine, methotrexate or phototherapy. Crohn's disease and Ulcerative colitis: Failure or clinically significant adverse effects to one of the formulary alternatives: budesonide, mesalamine or sulfasalazine.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **SYMDEKO**

### **Products Affected**

• SYMDEKO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of homozygous F508del mutation or at least one mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to tezacaftor/ivacaftor.
Age Restrictions	
Prescriber Restrictions	Infectious Disease specialist, Pulmonologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **SYMPAZAN**

### **Products Affected**

• SYMPAZAN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of difficulty or inability to swallow.
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to two of the formulary alternatives: carbamazepine, clonazepam, divalproex, ethosuximide, felbamate, gabapentin, lamotrigine, levetiracetam, lorazepam, oxcarbazepine, phenytoin, tiagabine, topiramate, valproic acid or zonisamide.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **SYNAREL**

### **Products Affected**

• SYNAREL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TACROLIMUS OINTMENT

### **Products Affected**

• tacrolimus topical

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to two of the topical formulary alternatives: clobetasol, betamethasone, fluocinolone, or fluocinonide.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **TAKHZYRO**

### **Products Affected**

• TAKHZYRO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Diagnosis of hereditary angioedema (HAE), must be confirmed by blood testing.
Age Restrictions	
Prescriber Restrictions	Allergist, Immunologist, Hematologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **TALTZ**

### **Products Affected**

• TALTZ AUTOINJECTOR

#### • TALTZ SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Plaque psoriasis: documentation of psoriasis of greater than 3% BSA or affecting crucial body areas such as hands, feet, face or genitals.
Age Restrictions	
Prescriber Restrictions	Dermatologist, Rheumatologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Psoriatic arthritis: Failure or clinically significant adverse effects to the formulary alternative: methotrexate. Ankylosing spondylitis: Failure or clinically significant adverse effects to one of the formulary alternatives: celecoxib, diclofenac, indomethacin, naproxen, or sulindac. Plaque psoriasis: Failure or clinically significant adverse effects to one of the formulary alternatives: acitretin, cyclosporine, methotrexate or phototherapy.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **TARGRETIN**

### **Products Affected**

• bexarotene

### • TARGRETIN TOPICAL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Dermatologist, Oncologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **TAZORAC**

#### **Products Affected**

- tazarotene topical cream
- tazarotene topical gel

- TAZORAC TOPICAL CREAM 0.05 % TAZORAC TOPICAL GEL

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Plaque psoriasis: Failure or clinically significant adverse effects to one of the topical formulary alternatives: calcipotriene, clobetasol or fluocinonide. Acne vulgaris: Failure or clinically significant adverse effects to two of the formulary alternatives: benzoyl peroxide/clindamycin topical, benzoyl peroxide/erythromycin topical, clindamycin topical, doxycycline oral, erythromycin topical, minocycline oral, tetracycline oral or tretinoin topical.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **TECFIDERA**

### **Products Affected**

• dimethyl fumarate

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to all of the formulary alternatives: Aubagio and glatiramer.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **TESTOSTERONE**

#### **Products Affected**

- methyltestosterone oral capsule
- testosterone cypionate intramuscular oil 100 mg/ml, 200 mg/ml
- testosterone enanthate

- testosterone transdermal gel in metereddose pump
- testosterone transdermal gel in packet
- testosterone transdermal solution in metered pump w/app

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documented pretreatment serum testosterone levels less than the laboratory's lower reference limit within the recent 3 months.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **TESTOSTERONE PUMP**

#### **Products Affected**

- testosterone transdermal gel in metered- testosterone transdermal gel in packet dose pump

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documented pretreatment serum testosterone levels less than the laboratory's lower reference limit within the recent 3 months.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to two of the formulary alternatives: testosterone cypionate, testosterone enanthate or testosterone transdermal gel or solution.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **TETRABENAZINE**

### **Products Affected**

• tetrabenazine

PA Criteria	Criteria Details
Exclusion Criteria	Concurrent use with an MAOI. Untreated or inadequately-treated depression, or current suicidality.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **TIGECYCLINE**

### **Products Affected**

• tigecycline

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Infectious Disease specialist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# TOBI PODHALER

### **Products Affected**

• TOBI PODHALER

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Infectious Disease specialist, Pulmonologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **TOBRAMYCIN SOLUTION**

### **Products Affected**

• tobramycin in 0.225 % nacl

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Infectious Disease specialist, Pulmonologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Subject to Part B vs Part D determination.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **TOLCAPONE**

### **Products Affected**

• tolcapone

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of concurrent use with levodopa and carbidopa.
Age Restrictions	
Prescriber Restrictions	Neurologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to two of the formulary alternatives: amantadine, bromocriptine, carbidopa/levodopa, entacapone, pramipexole, ropinirole, or selegiline.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **TRELSTAR**

### **Products Affected**

• TRELSTAR INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Subject to Part B vs Part D determination.
Indications	All Medically-accepted Indications.
Off Label Uses	
Part B Prerequisite	No

# **TRIENTINE**

### **Products Affected**

• trientine oral capsule 250 mg

PA Criteria	Criteria Details
Exclusion Criteria	Biliary cirrhosis, rheumatoid arthritis.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to the formulary alternative: penicillamine.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **TRIHEXYPHENIDYL**

### **Products Affected**

• trihexyphenidyl

PA Criteria	Criteria Details
Exclusion Criteria	Narrow angle glaucoma.
Required Medical Information	
Age Restrictions	Approve if less than 65 years of age. If aged 65 years and older, prior authorization criteria will apply.
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Parkinsonism: Failure or clinically significant adverse effects to one of the formulary alternatives: amantadine, bromocriptine, carbidopa/levodopa, entacapone, pramipexole, ropinirole, or selegiline. Medication-induced movement disorder - extrapyramidal disease: Failure or clinically significant adverse effects to the formulary alternative: amantadine.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **TRIMIPRAMINE**

### **Products Affected**

• trimipramine

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	Approve if less than 65 years of age. If aged 65 years and older, prior authorization criteria will apply.
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to two of the formulary alternatives: citalopram, escitalopram, fluoxetine, sertraline, venlafaxine, desvenlafaxine, duloxetine, mirtazapine, or bupropion
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **TRINTELLIX**

### **Products Affected**

• TRINTELLIX

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to two of the formulary alternatives: citalopram, escitalopram, fluoxetine, sertraline, venlafaxine, desvenlafaxine, duloxetine, mirtazapine, or bupropion
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **UBRELVY**

### **Products Affected**

• UBRELVY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Medical justification specifying that TWO formulary anti-migraine drugs from different classes have been tried and failed are contraindicated, or would not be medically appropriate. Classes include: (1) Analgesics-aspirin, naproxen, ibuprofen, diclofenac, celecoxib, indomethacin, nabumetone, and (2) Triptans- sumatriptan, rizatriptan/rizatriptan ODT
Age Restrictions	
Prescriber Restrictions	Neurologist, Pain Specialist, Headache Specialist
Coverage Duration	Until the end of calendar year
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **VALCHLOR**

### **Products Affected**

• VALCHLOR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Dermatologist, Oncologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **VANCOMYCIN CAPSULE**

### **Products Affected**

• vancomycin oral capsule 125 mg, 250 mg

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	C diff diarrhea: Reauthorization: Documentation of C. Difficile positive stool
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	3 months.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **VASCEPA**

### **Products Affected**

• icosapent ethyl oral capsule 0.5 gram, 1 gram

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Reduction of risk for myocardial infarction, stroke, coronary revascularization, and unstable angina requiring hospitalization: Documentation of maximally tolerated statin therapy, unless intolerant or contraindicated, with elevated triglycerides greater than or equal to 150 mg/dL and with either established cardiovascular disease or diabetes mellitus with two or more additional risk factors for cardiovascular disease. Severe hypertriglyceridemia: Documentation of elevated triglycerides greater than or equal to 500 mg/dL.
Age Restrictions	
Prescriber Restrictions	Lipidologist, Cardiologist, Endocrinologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to the formulary alternative: omega-3 acid ethyl esters.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **VEMLIDY**

### **Products Affected**

• VEMLIDY

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Hepatologist, Gastroenterologist, Infectious Disease specialist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **VERQUVO**

### **Products Affected**

• VERQUVO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Cardiologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **VIBERZI**

### **Products Affected**

• VIBERZI

PA Criteria	Criteria Details
Exclusion Criteria	History of gallbladder removal.
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to the all of the formulary alternatives: dicyclomine and loperamide.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **VIIBRYD**

#### **Products Affected**

- VIIBRYD ORAL TABLET
- VIIBRYD ORAL TABLETS,DOSE PACK 10 MG (7)- 20 MG (23)

• vilazodone

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to two of the formulary alternatives: citalopram, escitalopram, fluoxetine, sertraline, venlafaxine, desvenlafaxine, duloxetine, mirtazapine, or bupropion
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **VORICONAZOLE**

### **Products Affected**

• voriconazole intravenous

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **XATMEP**

### **Products Affected**

• XATMEP

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Oncologist, Pediatrician, Rheumatologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Subject to Part B vs Part D determination.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **XELJANZ**

### **Products Affected**

- XELJANZ ORAL SOLUTION
- XELJANZ ORAL TABLET

• XELJANZ XR

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Gastroenterologist, Rheumatologist
Coverage Duration	Until the end of calendar year.
Other Criteria	Rheumatoid Arthritis: Failure or clinically significant adverse effects to one of the formulary alternatives: azathioprine, cyclosporine, hydroxychloroquine, leflunomide, methotrexate, or sulfasalazine. Psoriatic arthritis: Failure or clinically significant adverse effects to the formulary alternative: methotrexate. Ulcerative colitis: Failure or clinically significant adverse effects to formulary alternative: Humira.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **XGEVA**

### **Products Affected**

• XGEVA

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Oncologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **XIFAXAN**

### **Products Affected**

• XIFAXAN

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Hepatic encephalopathy: Failure or clinically significant adverse effects to the formulary alternative: lactulose. Irritable bowel syndrome with diarrhea: Failure or clinically significant adverse effects to all of the formulary alternative: loperamide. Traveler's diarrhea: Failure or clinically significant adverse effects to one of the formulary alternatives: ciprofloxacin or levofloxacin.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

# **XOLAIR**

### **Products Affected**

• XOLAIR

PA Criteria	Criteria Details	
Exclusion Criteria		
Required Medical Information	Asthma (Initial): Forced expiratory volume in one second or peak expiratory flow less than or equal to 80% of predicted level, or measures of asthma control indicate uncontrolled asthma (eg, Asthma Control Test [ACT] score 19 or less). Baseline (pre-Xolair treatment) serum total IgE level greater than or equal to 30 IU/mL. Positive skin test or in vitro reactivity to a perennial aeroallergen.	
Age Restrictions		
Prescriber Restrictions	Allergist, Dermatologist, Immunologist, Pulmonologist, Otolaryngologist	
Coverage Duration	Until the end of calendar year.	
Other Criteria		
Indications	All FDA-approved Indications.	
Off Label Uses		
Part B Prerequisite	No	

## **XYREM**

### **Products Affected**

• sodium oxybate

### • XYREM

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Daytime excessive sleepiness in patients with narcolepsy: Failure or clinically significant adverse effects to two of the formulary alternatives: dextroamphetamine, methylphenidate, and modafinil.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## **ZARXIO**

### **Products Affected**

• ZARXIO

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	
Age Restrictions	
Prescriber Restrictions	Hematologist, Infectious Disease specialist, Oncologist
Coverage Duration	Until the end of calendar year.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

## ZEPATIER

### **Products Affected**

• ZEPATIER

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documentation of chronic hepatitis C infection confirmed by a detectable serum hepatitis C virus RNA through quantitative assay. Documentation of genotype. Documentation of the absence or presence of cirrhosis and if compensated or decompensated. Documentation of any previous treatment. Documentation of liver transplant status. Treatment regimen will be approved based on genotype and previous treatment experience as defined by current AASLD guidelines. For genotype 1a: Documentation for NS5A polymorphism testing.
Age Restrictions	
Prescriber Restrictions	Gastroenterologist, Hepatologist, Infectious Disease specialist
Coverage Duration	Duration will be applied consistent with current AASLD/IDSA guidance.
Other Criteria	
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### ZYPREXA RELPREVV

#### **Products Affected**

• ZYPREXA RELPREVV INTRAMUSCULAR SUSPENSION FOR RECONSTITUTION 210 MG

PA Criteria	Criteria Details
Exclusion Criteria	
Required Medical Information	Documented history of receiving oral olanzapine without any clinically significant side effects.
Age Restrictions	
Prescriber Restrictions	
Coverage Duration	Until the end of calendar year.
Other Criteria	Failure or clinically significant adverse effects to two of the formulary alternatives: Invega Sustenna, Invega Trinza or Risperdal Consta.
Indications	All FDA-approved Indications.
Off Label Uses	
Part B Prerequisite	No

### **INDEX**

A	AUBAGIO 22
ABELCET 1	AUGTYRO15, 17
ABILIFY ASIMTUFII	AUSTEDO ORAL TABLET 12 MG, 6 MG,
INTRAMUSCULAR	9 MG23
SUSPENSION, EXTENDED REL	AUVELITY 24
SYRING 720 MG/2.4 ML, 960 MG/3.2	AYVAKIT ORAL TABLET 100 MG, 200
ML2	MG, 25 MG, 300 MG, 50 MG 15, 17
ABILIFY MAINTENA	В
INTRAMUSCULAR	BALVERSA ORAL TABLET 3 MG, 4
SUSPENSION, EXTENDED REL	MG, 5 MG 15, 17
RECON 300 MG, 400 MG2	BESREMI 15, 17
ABILIFY MAINTENA	bexarotene
INTRAMUSCULAR	BOSULIF 15, 17
SUSPENSION, EXTENDED REL	BRAFTOVI
SYRING2	BRONCHITOL25
abiraterone oral tablet 250 mg, 500 mg 15,	BRUKINSA
17	C
acitretin	CABOMETYX 15, 17
ACTHAR 64	CALQUENCE (ACALABRUTINIB MAL)
ACTIMMUNE4	
adefovir5	CAPLYTA 107
AFINITOR DISPERZ ORAL TABLET	CAPRELSA 15, 17
FOR SUSPENSION 2 MG 15, 17	caspofungin26
AIMOVIG AUTOINJECTOR7	CIMZIA 27
AKEEGA 15, 17	CIMZIA POWDER FOR RECONST 27
ALECENSA	clobazam oral suspension 14
alosetron8	clobazam oral tablet14
ALUNBRIG 15, 17	clomipramine
alyq117	COMETRIQ 15, 17
AMBISOME9	COPIKTRA
ambrisentan 115	CORLANOR ORAL SOLUTION30
amitriptyline10	CORLANOR ORAL TABLET 30
amitriptyline-chlordiazepoxide oral tablet	COSENTYX (2 SYRINGES)31
12.5-5 mg, 25-10 mg	COSENTYX PEN (2 PENS) 31
amoxapine11	COSENTYX SUBCUTANEOUS
amphotericin b	SYRINGE 75 MG/0.5 ML31
amphotericin b liposome9	COSENTYX UNOREADY PEN 31
aprepitant18	COTELLIC 15, 17
APTIOM ORAL TABLET 200 MG, 400	cyclobenzaprine oral tablet 10 mg, 5 mg 32
MG, 600 MG, 800 MG13	D
ARCALYST 19	dalfampridine33
armodafinil20	DALIRESP34
asenapine maleate 107	DAURISMO ORAL TABLET 100 MG, 25
atovaquone21	MG15, 17

deferasirox oral tablet, dispersible 35	FETZIMA ORAL CAPSULE,EXT REL
desipramine36	24HR DOSE PACK 20 MG (2)- 40 MG
dexmethylphenidate oral tablet	(26)53
DIACOMIT14	FETZIMA ORAL CAPSULE,EXTENDED
diazepam intensol	RELEASE 24 HR 53
diazepam oral solution 5 mg/5 ml (1 mg/ml)	fingolimod58
	FINTEPLA14
dimethyl fumarate	FIRMAGON KIT W DILUENT SYRINGE
disopyramide phosphate oral capsule 39	
doxepin oral capsule40	FOTIVDA 15, 17
doxepin oral concentrate40	FRUZAQLA ORAL CAPSULE 1 MG, 5
DRIZALMA SPRINKLE 41	MG
dronabinol	FULPHILA118
droxidopa43	FYCOMPA ORAL SUSPENSION 13
DUAVEE 51	FYCOMPA ORAL TABLET13
DUPIXENT PEN 44, 45	G
DUPIXENT SYRINGE 44, 45	GAVRETO 15, 17
E	gefitinib
ELIGARD46	GENOTROPIN
ELIGARD (3 MONTH)46	GENOTROPIN MINIQUICK
ELIGARD (4 MONTH) 46	GILENYA
ELIGARD (6 MONTH)46	GILOTRIF
EMSAM	glatiramer
ENBREL 48	glatopa 60
ENBREL MINI 48	GLEOSTINE
ENBREL SURECLICK	glyburide
	e •
ENDARI	glyburide micronized
ERIVEDGE	glyburide-metformin oral tablet 1.25-250
	mg, 2.5-500 mg, 5-500 mg
ERLEADA ORAL TABLET 240 MG, 60	guanfacine oral tablet
MG	guanfacine oral tablet extended release 24 hr
erlotinib oral tablet 100 mg, 150 mg, 25 mg	63
	H
ESBRIET ORAL CAPSULE	HUMIRA PEN
ESBRIET ORAL TABLET 267 MG, 801	HUMIRA PEN CROHNS-UC-HS START
MG50	
estradiol oral	HUMIRA PEN PSOR-UVEITS-ADOL HS
estradiol transdermal patch weekly 51	68
everolimus (antineoplastic) oral tablet 15, 17	HUMIRA SUBCUTANEOUS SYRINGE
everolimus (antineoplastic) oral tablet for	KIT 40 MG/0.8 ML 66
suspension 3 mg, 5 mg 15, 17	HUMIRA(CF)66
EXKIVITY 15, 17	HUMIRA(CF) PEDI CROHNS STARTER
F	67
FANAPT	HUMIRA(CF) PEN66
fentanyl citrate buccal lozenge on a handle	HUMIRA(CF) PEN CROHNS-UC-HS 66
52	HUMIRA(CF) PEN PEDIATRIC UC 67

HUMIRA(CF) PEN PSOR-UV-ADOL HS	KORLYM 82
66	KOSELUGO ORAL CAPSULE 10 MG, 25
I	MG15, 17
IBRANCE	KRAZATI15, 17
icatibant54	L
ICLUSIG ORAL TABLET 10 MG, 15 MG,	lapatinib
30 MG, 45 MG 15, 17	LATUDA ORAL TABLET 120 MG, 20
icosapent ethyl oral capsule 0.5 gram, 1	MG, 40 MG, 60 MG, 80 MG 107
gram 174	ledipasvir-sofosbuvir
IDHIFA 15, 17	lenalidomide
imatinib oral tablet 100 mg, 400 mg 15, 17	LENVIMA ORAL CAPSULE 10 MG/DAY
IMBRUVICA ORAL CAPSULE 140 MG,	(10 MG X 1), 12 MG/DAY (4 MG X 3),
70 MG	14 MG/DAY(10 MG X 1-4 MG X 1), 18
IMBRUVICA ORAL SUSPENSION. 15, 17	MG/DAY (10 MG X 1-4 MG X2), 20
IMBRUVICA ORAL TABLET 140 MG,	MG/DAY (10 MG X 2), 24 MG/DAY(10
280 MG, 420 MG, 560 MG 15, 17	MG X 2-4 MG X 1), 4 MG, 8 MG/DAY
imipramine hel	(4 MG X 2) 15, 17
imipramine pamoate	LEUKINE INJECTION RECON SOLN 84
INCRELEX70	leuprolide (3 month) 85
indomethacin oral capsule71	leuprolide subcutaneous kit
INGREZZA72	levalbuterol hel inhalation solution for
INGREZZA INITIATION PK(TARDIV) 72	nebulization 0.31 mg/3 ml, 0.63 mg/3 ml,
INGREZZA SPRINKLE	1.25 mg/0.5 ml, 1.25 mg/3 ml 86
INLYTA ORAL TABLET 1 MG, 5 MG. 15,	LIBERVANT97
17	lidocaine topical adhesive patch, medicated 5
INQOVI 15, 17	%87
INREBIC	LONSURF 15, 17
INVEGA HAFYERA74	LORBRENA ORAL TABLET 100 MG, 25
INVEGA SUSTENNA 75	MG15, 17
INVEGA TRINZA76	LUMAKRAS ORAL TABLET 120 MG,
IRESSA	320 MG
itraconazole oral solution	LUPRON DEPOT88
IWILFIN 15, 17	LUPRON DEPOT (3 MONTH) 88
J	LUPRON DEPOT (4 MONTH) 88
JAKAFI	LUPRON DEPOT (6 MONTH) 88
JAYPIRCA ORAL TABLET 100 MG, 50	lurasidone oral tablet 120 mg, 20 mg, 40
MG	mg, 60 mg, 80 mg107
JYLAMVO	LYNPARZA
K	LYTGOBI 12 MG DOSE (3X 4 MG TB) 15,
KALYDECO79	17
KERENDIA	LYTGOBI 16 MG DOSE (4X 4 MG TB) 15,
KINERET	17
KISQALI FEMARA CO-PACK 15, 17	LYTGOBI 20 MG DOSE (5X 4 MG TB) 15,
KISQALI ORAL TABLET 200 MG/DAY	17
(200 MG X 1), 400 MG/DAY (200 MG X	-1
2), 600 MG/DAY (200 MG X 3) 15, 17	

methylphenidate hcl oral tablet extended release 6 methyltestosterone oral capsule 159 metryrosine 95 modafinil 96 molindone oral tablet 10 mg, 25 mg, 5 mg65 MOTPOLY XR ORAL CAPSULE, EXTENDED RELEASE 24HR 100 MG, 150 MG, 200 MG 14 MOVANTIK 29 NAYZILAM 97 OTEZLA STARTER ORAL TABLET ORAL TABLET S,DOSE PACK 10 MG (4)-20 MG (4)-30 MG (47) 110 oxandrolone 111 OXBRYTA ORAL TABLET 300 MG, 500 MG (4)-30 MG (47) 110 OXBRYTA ORAL TABLET 300 MG, 500 MG 111 OXBRYTA ORAL TABLET FOR SUSPENSION 112 OXERVATE 113 OXTELLAR XR ORAL TABLET EXTENDED RELEASE 24 HR 150 MG, 300 MG, 600 MG 114 OZEMPIC SUBCUTANEOUS PEN	LYTGOBI ORAL TABLET 4 MG, 4 MG (4X 4 MG TB), 4 MG (5X 4 MG TB). 16,	ODOMZO
MATULANE         16, 17         megestrol oral suspension 400 mg/10 ml (40 mg/ml)         MCRESTOR MORESTITUTION         16, 17           MEKINIST ORAL RECON SOLN         16, 17         MG. 200 MG         16, 17           MEKINIST ORAL TABLET 0.5 MG, 2         MG         16, 17           MEKTOVI         16, 17         MC G. 200 MG         16, 17           MEKTOVI         16, 17         MC G. 200 MG         16, 17           MEKTOVI         16, 17         MC G. 200 MG         16, 17           MEKTOVI         16, 17         MC G. 200 MG         16, 17           MEKTOVI         16, 17         MC G. 200 MG         16, 17           MEKTOVI         16, 17         MC G. 200 MG         16           MEXTOVI         16, 17         MC G. 200 MG         16           MEXTOVI         16, 17         MC G. 200 MG         16           MELONARIA SUSPENSION TOR         10         ONUREG         16, 17           MELONARIA SUSPENSION TOR         10         ONUREG         106           MELONARIA SUSPENSION TOR         125         MG/M MILOLA M		
megestrol oral suspension 400 mg/10 ml (40 mg/ml)         RECONSTITUTION         16, 17           megestrol oral tablet         90         MEKINIST ORAL RECON SOLN         16, 17           MEKINIST ORAL TABLET 0.5 MG, 2         0         MG, 200 MG         16, 17           MEKINIST ORAL TABLET 0.5 MG, 2         16, 17         MG, 200 MG         16, 17           MEKTOVI         16, 17         olanzapine oral tablet, disintegrating         104           methovasalen         91         MG         200 MG         16, 17           methoxsalen         93         ONUREG         16, 17           methylphenidate hel oral capsule, er         94         MG/0.7 ML         108           methylphenidate hel oral capsule, er         66         MG/0.7 ML         108           biphasis 30-70         6         GROVYX         16, 17           methylphenidate hel oral tablet         6         MG         MG/0.7 ML         108           MG 2         MG/0.7 ML         108         MG/0.7 ML         10           MEXAMBI         109         ORSERDU ORAL TABLET 345 MG, 86         MG         MG/0.7 ML         10           MEXAMBI         10 oran         MG/0.7 ML         10         MCRAMBI         10           MEXAMBI         1		
mg/ml)         90         OJEMDA ORAL TABLET         16, 17           MEKINIST ORAL RECON SOLN         16, 17         MG (20) MG         16, 17           MEKINIST ORAL TABLET 0.5 MG, 2         MG         16, 17           MEKTOVI         16, 17         olanzapine intramuscular         105           methocarbamol oral         92         olanzapine oral tablet, disintegrating         104           methoxasalen         91         oNUREG         16, 17           methologa- hydrochlorothiazide         94         oRENCIA CLICKJECT         108           methylphenidate hel oral capsule, er biphasic         oBoylog         16, 17         oRENCIA SUBCUTANEOUS SYRINGE         125 MG/ML, 50 MG/0.4 ML, 87.5         MG/0.7 ML         108           methylphenidate hel oral capsule, er biphasic         oBoylog         100         ORSERDU ORAL TABLET 100 MG, 150 MG, 00         100         ORDIVEG         104         MNITROPE         106         ONUREG         106         ORENCIA SUBCUTANEOUS SYRINGE         125 MG/ML, 50 MG/0.4 ML, 87.5         MG/0.7 ML         108         ORGOVYX         16, 17         ORKAMBI         109         ORSERDU ORAL TABLET 345 MG, 86         MG         107         ORSERDU ORAL TABLET 100 MG, 100         MG/0.7 ML         108         ORGOVYX         16, 17         ORSERDU ORAL TABLET 100 MG, 100	magastrol oral suspension 400 mg/10 ml (40	
megestrol oral tablet         .90           MEKINIST ORAL RECON SOLN         .16, 17           MEKINIST ORAL TABLET 0.5 MG, 2         .90           MG         .16, 17           MEKINIST ORAL TABLET 0.5 MG, 2         .10           MEKTOVI         .16, 17           methopoarbanate         .91           methooxalen         .92           methoxsalen         .93           methyladpa-hydrochlorothiazide         .94           methylphenidate hel oral capsule, er biphasic 30-70         .6           biphasic 30-70         .6           methylphenidate hel oral capsule, er biphasic 50-50         .6           methylphenidate hel oral solution         .6           methylphenidate hel oral tablet         .6           methylphenidate hel oral tablet extended release         .6           methylphenidate hel oral tablet extended release         .95           modafinil         .95           modafinil         .96           molindone oral tablet 10 mg, 25 mg, 5 mg65           MOTPOLY XR ORAL         .9           CAPSULE,EXTENDED RELEASE         .24HR 100 MG, 150 MG, 200 MG           24HR 100 MG, 150 MG, 200 MG         .10           NYZILAM         .97           NERLYNX         .16, 17		· · · · · · · · · · · · · · · · · · ·
MEKINIST ORAL RECON SOLN 16, 17         MG, 200 MG         16, 17           MEKINIST ORAL TABLET 0.5 MG, 2         10, 17           MG         16, 17           MEKTOVI         16, 17           methopobamate         91           methooxalen         92           methoyldopa-hydrochlorothiazide         94           methylphenidate hel oral capsule, er biphasic 30-70         6           methylphenidate hel oral capsule, er biphasic 30-70         6           methylphenidate hel oral capsule, er biphasic 30-70         6           methylphenidate hel oral solution         6           methylphenidate hel oral sublet         6           methylphenidate hel oral tablet extended release         6           methylptestosterone oral capsule         159            mettyrosine         95           modafinil         96           molindone oral tablet 10 mg, 25 mg, 5 mg65         MG (4)-30 MG (47)         110           MG (2)-30 MG, 500 MG         11           MG (4)-30 MG, 500 MG         10           MG (4)-30 MG (47)         110           OxBRYTA ORAL TABLET 300 MG, 500         112           MG (4)-30 MG (47)         110           OxBRYTA ORAL TABLET 300 MG, 500         11           MG (5)-30		
MEKINIST ORAL TABLET 0.5 MG, 2         olanzapine intramuscular         105           MEKTOVI         16, 17         OMNITROPE         106           methocarbamol oral         92         ONUREG         16, 17           methoxalen         93         ORENCIA CLICKJECT         108           methyldopa-hydrochlorothiazide         94         MG/0.7 ML         108           methylphenidate hcl oral capsule, er biphasic         50-50         6         MG/0.7 ML         108           50-50         6         MG/0.7 ML         108         MG/0.7 ML         108           methylphenidate hcl oral capsule, er biphasic         6         MG/0.7 ML         108         MG/0.7 ML         108           50-50         6         MGKAMBI         109         MG/0.7 ML         108           Methylphenidate hcl oral tablet         6         MG/0.7 ML         108         MG/0.7 ML         108           methylphenidate hcl oral tablet extended release         6         MG/0.7 ML         10         MG/0.7 ML         10         MC         MC         MC         11         OTEZLA         110         OTEZLA STARTER ORAL         TABLETS,DOSE PACK 10 MG (4)-20         MG/0.1 MG (4)-20         MG/0.1 MG (4)-30         MG/0.1 MG (4)-20         MG/0.1 MG (4)-30         MG/0.1		
MG         16, 17         olanzapine oral tablet, disintegrating         104           MEKTOVI         16, 17         oMNTROPE         106           merprobamate         91         oMNTROPE         106           methocarbamol oral         92         oRENCIA CLICKJECT         108           methoxsalen         93         oRENCIA SUBCUTANEOUS SYRINGE         125 MG/ML, 50 MG/0.4 ML, 87.5           methylphenidate hel oral capsule, er biphasic         oRGOVYX         16, 17           50-50         6         oRGOVYX         16, 17           methylphenidate hel oral solution         6         oRGOVYX         16, 17           methylphenidate hel oral solution         6         oRGOVYX         16, 17           methylphenidate hel oral tablet         6         oRGOVYX         16, 17           methylphenidate hel oral tablet         6         MG         110           methylphenidate hel oral tablet         6         MG         111           methylptesiotserone oral capsule         159         methylitestosterone oral tablet 300 mg         82           methylitestosterone oral tablet 10 mg, 25 mg, 5 mg65         MG         (4)-30 MG (47)         110           OXBRYTA ORAL TABLET 300 MG, 500         MG         112           CAPSULE, EXTEN		
MEKTOVI.         16, 17         OMNITROPE         106           metporbamate         91         ONUREG         16, 17           methocarbamol oral         92         ORENCIA CLICKJECT         108           methyldopa-hydrochlorothiazide         94         ORENCIA SUBCUTANEOUS SYRINGE           methylphenidate hcl oral capsule, er         biphasic 30-70         6           methylphenidate hcl oral capsule, er biphasic         ORGOVYX         16, 17           50-50         6         ORKAMBL         109           ORSERDU ORAL TABLET 345 MG, 86         MG         16, 17           methylphenidate hcl oral solution         6         MG         16, 17           methylphenidate hcl oral solution         6         MG         16, 17           methylphenidate hcl oral solution         6         MG         16, 17           methylphenidate hcl oral tablet         6         MG         16, 17           mettyrosine         95         mctyrosine         95           molafinil         96         MG (4)-30 MG (47)         110           MOTPOLY XR ORAL         20         XBRYTA ORAL TABLET 300 MG, 500           MOTPOLY XR ORAL         29         XBRYTA ORAL TABLET FOR         SUSPENSION         112           NAY	· · · · · · · · · · · · · · · · · · ·	
Methocarbamol oral		
methocarbamol oral         92         ORENCIA CLICKJECT         108           methyldopa-hydrochlorothiazide         94         ORENCIA SUBCUTANEOUS SYRINGE           methylphenidate hcl oral capsule, er biphasic 30-70         6         MG/0.7 ML         108           50-50         6         MG/0.7 ML         108           methylphenidate hcl oral capsule, er biphasic 50-50         6         MG/0.7 ML         109           50-50         6         MG/0.7 ML         109           methylphenidate hcl oral solution         6         MG         16, 17           methylphenidate hcl oral tablet         6         MG         16, 17           methylphenidate hcl oral tablet extended release         6         MG         110           methylphenidate hcl oral tablet extended release         0         MG         110           methylphenidate hcl oral tablet extended release         0         MG         110           methylphenidate hcl oral tablet oral tab		
methoxsalen         93           methyldopa-hydrochlorothiazide         94           methylphenidate hol oral capsule, er biphasic 30-70         6           methylphenidate hol oral capsule, er biphasic 50-50         6           methylphenidate hol oral capsule, er biphasic 50-50         0RGOVYX         16, 17           omethylphenidate hol oral solution         6         0RGOVYX         16, 17           methylphenidate hol oral solution         6         MG         16, 17           methylphenidate hol oral tablet ablet oral tablet extended release         6         MG         16, 17           methylphenidate hol oral tablet extended release         6         MG         10         0TEZLA         110           methylphenidate hol oral tablet extended release         6         MG         10         0TEZLA         110           methylphenidate hol oral tablet extended release         6         MG         10         0TEZLA         110           methylphenidate hol oral tablet extended release         6         MG         40		
methyldopa-hydrochlorothiazide         94           methylphenidate hel oral capsule, er         biphasic 30-70         6           methylphenidate hel oral capsule, er biphasic         6         ORGOVYX         16, 17           50-50         6         ORKAMBI         109           methylphenidate hel oral solution         6         MG         16, 17           methylphenidate hel oral tablet         6         MG         16, 17           methylphenidate hel oral tablet extended release         6         MG         16, 17           methyltestosterone oral capsule         159         MG (4)-30 MG (47)         110           methyltestosterone oral tablet 10 mg, 25 mg, 5 mg65         MG         10         XBRYTA ORAL TABLET 300 MG, 500           MOTPOLY XR ORAL         NG         112         OXBRYTA ORAL TABLET 300 MG, 500           MG         112         OXBRYTA ORAL TABLET TOR         SUSPENSION         112           MOVANTIK         29         N         SUSPENSION         112           MOVANTIK         97         NERLARX         16, 17         NINEATO         16, 17           NINLARO         16, 17         NINLARO         16, 17         NINLARO         16, 17           NUEDEXTA         100         PANRETIN		
methylphenidate hel oral capsule, er biphasic 30-70         6         MG/0.7 ML         108           50-50         6         ORGOVYX         16, 17           methylphenidate hel oral capsule, er biphasic 50-50         6         ORKAMBI         109           methylphenidate hel oral solution         6         ORSERDU ORAL TABLET 345 MG, 86         MG         16, 17           methylphenidate hel oral tablet         6         MG         16, 17         OTEZLA         110           methylphenidate hel oral tablet extended release         6         MG         16, 17         OTEZLA         110           methylptestosterone oral capsule         159         MG (4)-30 MG (47)         110         OXBRYTA ORAL TABLET SOM MG, 500         MG         111           motifepristone oral tablet 10 mg, 25 mg, 5 mg65         MOTPOLY XR ORAL         SUSPENSION         112           CAPSULE,EXTENDED RELEASE         24HR 100 MG, 150 MG, 200 MG         14         OXERVATE         113           OXFELLAR XR ORAL TABLET         OXERVATE         113         OXTELLAR XR ORAL TABLET           MOVANTIK         29         N         N         300 MG, 600 MG         114           NAYZILAM         97         NIJECTOR 0.25 MG OR 0.5 MG (2         MG/3 ML), 0.25 MG OR 0.5 MG (2           NIVESTYM INJECTI		
biphasic 30-70		
methylphenidate hcl oral capsule,er biphasic         ORKAMBI         109           50-50         6         MG         16, 17           methylphenidate hcl oral solution         6         MG         16, 17           methylphenidate hcl oral tablet         6         MG         16, 17           methylphenidate hcl oral tablet extended release         6         MG         10           methylphenidate hcl oral tablet extended release         6         MG         10           methylphenidate hcl oral tablet extended release         6         MG         10           methylphenidate hcl oral tablet extended release         6         MG         10           methylphenidate hcl oral tablet extended release         6         MG         11           methylphenidate hcl oral tablet extended release         6         MG         10           methylphenidate hcl oral tablet actended release         6         MG         10           methylphenidate hcl oral tablet actended release         6         MG         10           methylphenidate hcl oral tablet actended release         6         MG         10           methylphenidate hcl oral tablet actended release         6         MG         10           methylphenidate hcl oral tablet actended mclate for a tablet actended release		
50-50         6         MCRSERDU ORAL TABLET 345 MG, 86           methylphenidate hel oral solution         6         MG         16, 17           methylphenidate hel oral tablet         6         MG         16, 17           methylphenidate hel oral tablet extended release         6         TABLETS, DOSE PACK 10 MG (4)-20           methyltestosterone oral capsule         159         MG (4)-30 MG (47)         110           methyltestosterone oral tablet 300 mg         82         MG (4)-30 MG (47)         110           modafinil         96         MG (4)-30 MG (47)         110           modafinil         96         MG (4)-30 MG (47)         110           molindone oral tablet 10 mg, 25 mg, 5 mg65         MG         MG         11           MOTPOLY XR ORAL         SUSPENSION         112           OXERVATE         113         OXERVATE         113           OXEVATE         113         OXELLAR XR ORAL TABLET FOR         SUSPENSION         112           MOVANTIK         29         OXERVATE         113         OXELLAR XR ORAL TABLET         12           NAYZILAM         97         NERLYNX         16, 17         NUSECTOR 0.25 MG OR 0.5 MG (2         MG/3 ML), 0.25 MG OR 0.5 MG (2           MEXAVAR         16, 17         MG/1.5 ML), 1 MG		
methylphenidate hcl oral solution       6         methylphenidate hcl oral tablet       6         methylphenidate hcl oral tablet extended release       6         methyltestosterone oral capsule       159         metyrosine       95         modafinil       96         molindone oral tablet 10 mg, 25 mg, 5 mg65       MG         MOTPOLY XR ORAL       SUSPENSION         CAPSULE,EXTENDED RELEASE       OXERVATE         24HR 100 MG, 150 MG, 200 MG       14         MOVANTIK       29         N       OXERVATE         NEXIVA       16, 17         NINLARO       16, 17         NIVESTYM INJECTION       98         NOXAFIL ORAL SUSPENSION       121         NOXAFIL ORAL SUSPENSION       121         NUGLALA       99         NUEDEXTA       100         NUPLAZID       101         O       101         Pemenbarbital       16, 17         NUPLAZID       101         phenobarbital       119	· · · · · · · · · · · · · · · · · · ·	
methylphenidate hcl oral tablet         6           methylphenidate hcl oral tablet extended release         6           methyltestosterone oral capsule         159           modafini         96           molindone oral tablet 300 mg         82           modafini         96           molindone oral tablet 10 mg, 25 mg, 5 mg65         MG (4)-30 MG (47)           MG (24)-30 MG (47)         110           OXBRYTA ORAL TABLET 300 MG, 500         MG           MSRYTA ORAL TABLET FOR         SUSPENSION           SUSPENSION         112           OXERVATE         113           OXTELLAR XR ORAL TABLET         EXTENDED RELEASE 24 HR 150 MG, 300 MG, 600 MG           NAYZILAM         97           NEXAVAR         16, 17           NINLARO         16, 17		
methylphenidate hcl oral tablet extended release         OTEZLA STARTER ORAL           methyltestosterone oral capsule         159           methyltestosterone oral tablet 300 mg         20           modafinil         96           modafinil         96           Modafinil         96           Modafinil         96           Mod McJ Albert         112           OXBRYTA ORAL TABLET         20           OXERVATE         113           OXTELLAR XR ORAL TABLET         12           OXEMPIC SUBCUTANEOUS PEN         114           OZEMPIC SU		
release         6         TABLETS, DOSE PACK 10 MG (4)-20           methyltestosterone oral capsule         159         MG (4)-30 MG (47)         110           metyrosine         95         MG (4)-30 MG (47)         110           mifepristone oral tablet 300 mg         82         MG (4)-30 MG (47)         110           modafinil         96         MG (4)-30 MG (47)         110           modafinil         98         MG (4)-30 MG (47)         110           modafinil         98         MG (M) MG (M)         112           MD MC (A) Table Trick         99         MG (M) MG (M)         100           MD MC (A) Table Trick         112         OXBRYTA ORAL Table Trick         112           OXERVATE         113         OXTELLAR XR ORAL Table Trick         113           OXEVATE         113         OXTELLAR XR ORAL Table Trick         12           MAYZILAM         97         NEXAVAR         16,17	methylphenidate hcl oral tablet6	
methyltestosterone oral capsule       159       MG (4)-30 MG (47)       110         metyrosine       95       oxandrolone       111         mifepristone oral tablet 300 mg       82       OXBRYTA ORAL TABLET 300 MG, 500         modafinil       96       MG       112         molindone oral tablet 10 mg, 25 mg, 5 mg65       MG       112         MOTPOLY XR ORAL       SUSPENSION       112         CAPSULE,EXTENDED RELEASE       OXERVATE       113         24HR 100 MG, 150 MG, 200 MG       14       OXTELLAR XR ORAL TABLET         MOVANTIK       29       EXTENDED RELEASE 24 HR 150 MG,         N       300 MG, 600 MG       114         OZEMPIC SUBCUTANEOUS PEN       INJECTOR 0.25 MG OR 0.5 MG (2         MEXAVAR       16, 17       MG/1.5 ML), 1 MG/DOSE (4 MG/3         NIVESTYM INJECTION       98       ML), 2 MG/DOSE (8 MG/3 ML)       61         NUCALA       99       PANRETIN       16, 17         NUCALA       99       PANRETIN       16, 17         NUEDEXTA       100       PEMAZYRE       16, 17         NUPLAZID       101       perphenazine-amitriptyline       10         O       phenobarbital       119	methylphenidate hcl oral tablet extended	
metyrosine       95         mifepristone oral tablet 300 mg       82         modafinil       96         molindone oral tablet 10 mg, 25 mg, 5 mg65       MG         MOTPOLY XR ORAL       SUSPENSION         CAPSULE,EXTENDED RELEASE       OXERVATE         24HR 100 MG, 150 MG, 200 MG       14         MOVANTIK       29         N       SUSPENSION         NERLYNX       16, 17         NERLYNX       16, 17         NINLARO       16, 17         NIVESTYM INJECTION       98         NOXAFIL ORAL SUSPENSION       121         NUEDEAA       16, 17         NUEDEXTA       100         NUPLAZID       101         O       Demodafinil         95       MG         0XBRYTA ORAL TABLET FOR         SUSPENSION       112         OXTELLAR XR ORAL TABLET       113         OXEMPIC SUBCUTANEOUS PEN       INJECTOR 0.25 MG OR 0.5 MG (2         MG/3 ML), 0.25 MG OR 0.5 MG (2       MG/3 ML), 0.25 MG OR 0.5 MG (2         MG/3 ML), 1 MG/DOSE (8 MG/3 ML)       61         PANRETIN       16, 17         PANRETIN       16, 17         PEMAZYRE       16, 17         PEMAZYRE </td <td>release6</td> <td>TABLETS, DOSE PACK 10 MG (4)-20</td>	release6	TABLETS, DOSE PACK 10 MG (4)-20
mifepristone oral tablet 300 mg       82       OXBRYTA ORAL TABLET 300 MG, 500 MG         modafinil       96       MG       112         molindone oral tablet 10 mg, 25 mg, 5 mg65       MG       112         MOTPOLY XR ORAL       SUSPENSION       112         CAPSULE,EXTENDED RELEASE       OXERVATE       113         24HR 100 MG, 150 MG, 200 MG       14       OXERVATE       113         MOVANTIK       29       EXTENDED RELEASE 24 HR 150 MG, 300 MG, 600 MG       114         NAYZILAM       97       OZEMPIC SUBCUTANEOUS PEN         NERLYNX       16, 17       INJECTOR 0.25 MG OR 0.5 MG (2         NEXAVAR       16, 17       MG/3 ML), 0.25 MG OR 0.5 MG (2         NINLARO       16, 17       MG/1.5 ML), 1 MG/DOSE (4 MG/3         NIVESTYM INJECTION       98       ML), 2 MG/DOSE (8 MG/3 ML)       61         NUBEQA       16, 17       PANRETIN       16, 17         NUCALA       99       PANRETIN       16, 17         NUEDEXTA       100       PEMAZYRE       16, 17         NUPLAZID       101       perphenazine-amitriptyline       10         OXBRYTA ORAL TABLET 300 MG, 500 MG       112       OXERVATE       113         OXERVATE       NG       NG       00	methyltestosterone oral capsule 159	MG (4)-30 MG (47)110
modafinil       96       MG       112         molindone oral tablet 10 mg, 25 mg, 5 mg65       OXBRYTA ORAL TABLET FOR         MOTPOLY XR ORAL       SUSPENSION       112         CAPSULE, EXTENDED RELEASE       OXERVATE       113         24HR 100 MG, 150 MG, 200 MG       14       OXTELLAR XR ORAL TABLET         MOVANTIK       29       EXTENDED RELEASE 24 HR 150 MG, 300 MG, 600 MG       114         NAYZILAM       97       OZEMPIC SUBCUTANEOUS PEN         NERLYNX       16, 17       INJECTOR 0.25 MG OR 0.5 MG (2         NEXAVAR       16, 17       MG/3 ML), 0.25 MG OR 0.5 MG (2         NINLARO       16, 17       MG/1.5 ML), 1 MG/DOSE (4 MG/3         NIVESTYM INJECTION       98       ML), 2 MG/DOSE (8 MG/3 ML)       61         NUBEQA       16, 17       PANRETIN       16, 17         NUCALA       99       PANRETIN       16, 17         NUEDEXTA       100       PEMAZYRE       16, 17         NUPLAZID       101       perphenazine-amitriptyline       10         O       phenobarbital       119	metyrosine95	oxandrolone111
molindone oral tablet 10 mg, 25 mg, 5 mg65         OXBRYTA ORAL TABLET FOR SUSPENSION         112           MOTPOLY XR ORAL         SUSPENSION         112           CAPSULE,EXTENDED RELEASE         OXERVATE         113           24HR 100 MG, 150 MG, 200 MG         14         OXTELLAR XR ORAL TABLET           MOVANTIK         29         EXTENDED RELEASE 24 HR 150 MG, 300 MG, 600 MG         114           NAYZILAM         97         OZEMPIC SUBCUTANEOUS PEN           NERLYNX         16, 17         INJECTOR 0.25 MG OR 0.5 MG (2           NEXAVAR         16, 17         MG/3 ML), 0.25 MG OR 0.5 MG (2           NIVESTYM INJECTION         98         ML), 2 MG/DOSE (8 MG/3 ML)         61           NUCALA         99         PANRETIN         16, 17           NUCALA         99         PANRETIN         16, 17           NUPLAZID         101         PEMAZYRE         16, 17           NUPLAZID         101         perphenazine-amitriptyline         10           O         phenobarbital         119	mifepristone oral tablet 300 mg 82	OXBRYTA ORAL TABLET 300 MG, 500
MOTPOLY XR ORAL       SUSPENSION       112         CAPSULE,EXTENDED RELEASE       OXERVATE       113         24HR 100 MG, 150 MG, 200 MG       14       OXTELLAR XR ORAL TABLET         MOVANTIK       29       EXTENDED RELEASE 24 HR 150 MG,         N       300 MG, 600 MG       114         NAYZILAM       97       OZEMPIC SUBCUTANEOUS PEN         NERLYNX       16, 17       INJECTOR 0.25 MG OR 0.5 MG (2         NEXAVAR       16, 17       MG/3 ML), 0.25 MG OR 0.5 MG (2         NINLARO       16, 17       MG/1.5 ML), 1 MG/DOSE (4 MG/3         NIVESTYM INJECTION       98       ML), 2 MG/DOSE (8 MG/3 ML)       61         NUCALA       99       PANRETIN       16, 17         NUCALA       99       PANRETIN       16, 17         NUPLAZID       101       PEMAZYRE       16, 17         NUPLAZID       101       perphenazine-amitriptyline       10         O       phenobarbital       119	modafinil96	MG112
CAPSULE,EXTENDED RELEASE       OXERVATE       113         24HR 100 MG, 150 MG, 200 MG       14       OXTELLAR XR ORAL TABLET         MOVANTIK       29       EXTENDED RELEASE 24 HR 150 MG,         N       300 MG, 600 MG       114         NAYZILAM       97       OZEMPIC SUBCUTANEOUS PEN         NERLYNX       16, 17       INJECTOR 0.25 MG OR 0.5 MG (2         NEXAVAR       16, 17       MG/3 ML), 0.25 MG OR 0.5 MG (2         NINLARO       16, 17       MG/1.5 ML), 1 MG/DOSE (4 MG/3         NIVESTYM INJECTION       98       ML), 2 MG/DOSE (8 MG/3 ML)       61         NOXAFIL ORAL SUSPENSION       121       P         NUEDEATA       100       PANRETIN       16, 17         NUPLAZID       101       PEMAZYRE       16, 17         PEMAZYRE       16, 17       perphenazine-amitriptyline       10         Phenobarbital       119	molindone oral tablet 10 mg, 25 mg, 5 mg65	OXBRYTA ORAL TABLET FOR
24HR 100 MG, 150 MG, 200 MG       14       OXTELLAR XR ORAL TABLET         MOVANTIK       29       EXTENDED RELEASE 24 HR 150 MG,         N       300 MG, 600 MG       114         NAYZILAM       97       OZEMPIC SUBCUTANEOUS PEN         NERLYNX       16, 17       INJECTOR 0.25 MG OR 0.5 MG (2         NEXAVAR       16, 17       MG/3 ML), 0.25 MG OR 0.5 MG (2         NINLARO       16, 17       MG/1.5 ML), 1 MG/DOSE (4 MG/3         NIVESTYM INJECTION       98       ML), 2 MG/DOSE (8 MG/3 ML)       61         NOXAFIL ORAL SUSPENSION       121       P         NUCALA       99       PANRETIN       16, 17         NUCALA       99       PANRETIN       16, 17         NUPLAZID       101       PEMAZYRE       16, 17         PEMAZYRE       16, 17       perphenazine-amitriptyline       10         Phenobarbital       119	MOTPOLY XR ORAL	SUSPENSION 112
MOVANTIK       29       EXTENDED RELEASE 24 HR 150 MG, 300 MG, 600 MG.       114         NAYZILAM       97       OZEMPIC SUBCUTANEOUS PEN         NERLYNX       16, 17       INJECTOR 0.25 MG OR 0.5 MG (2         NEXAVAR       16, 17       MG/3 ML), 0.25 MG OR 0.5 MG (2         NINLARO       16, 17       MG/1.5 ML), 1 MG/DOSE (4 MG/3 ML)         NIVESTYM INJECTION       98       ML), 2 MG/DOSE (8 MG/3 ML)       61         NOXAFIL ORAL SUSPENSION       121       P         NUBEQA       16, 17       PANRETIN       16, 17         NUCALA       99       PANRETIN       16, 17         NUEDEXTA       100       PEMAZYRE       16, 17         NUPLAZID       101       Pemazyre       16, 17         Phenobarbital       119	CAPSULE, EXTENDED RELEASE	OXERVATE113
N       300 MG, 600 MG	24HR 100 MG, 150 MG, 200 MG 14	OXTELLAR XR ORAL TABLET
NAYZILAM       97         NERLYNX       16, 17         NEXAVAR       16, 17         NINLARO       16, 17         NIVESTYM INJECTION       98         NOXAFIL ORAL SUSPENSION       121         NUBEQA       16, 17         NUCALA       99         NUEDEXTA       100         NUPLAZID       101         O       O         OZEMPIC SUBCUTANEOUS PEN         INJECTOR 0.25 MG OR 0.5 MG(2         MG/3 ML), 0.25 MG OR 0.5 MG(2         MG/1.5 ML), 1 MG/DOSE (4 MG/3         ML), 2 MG/DOSE (8 MG/3 ML)       61         PANRETIN       16, 17         perphenazine-amitriptyline       16, 17         perphenazine-amitriptyline       10         phenobarbital       119	MOVANTIK29	EXTENDED RELEASE 24 HR 150 MG,
NERLYNX       16, 17       INJECTOR 0.25 MG OR 0.5 MG (2         NEXAVAR       16, 17       MG/3 ML), 0.25 MG OR 0.5 MG (2         NINLARO       16, 17       MG/1.5 ML), 1 MG/DOSE (4 MG/3         NIVESTYM INJECTION       98       ML), 2 MG/DOSE (8 MG/3 ML)       61         NOXAFIL ORAL SUSPENSION       121       P         NUBEQA       16, 17       PANRETIN       16, 17         NUCALA       99       pazopanib       16, 17         NUEDEXTA       100       PEMAZYRE       16, 17         NUPLAZID       101       perphenazine-amitriptyline       10         O       phenobarbital       119	N	300 MG, 600 MG114
NEXAVAR       16, 17         NINLARO       16, 17         NIVESTYM INJECTION       98         NOXAFIL ORAL SUSPENSION       121         NUBEQA       16, 17         NUCALA       99         NUEDEXTA       100         NUPLAZID       101         O       MG/3 ML), 0.25 MG OR 0.5 MG(2         MG/1.5 ML), 1 MG/DOSE (4 MG/3         ML), 2 MG/DOSE (8 MG/3 ML)         ML), 2	NAYZILAM 97	OZEMPIC SUBCUTANEOUS PEN
NINLARO	NERLYNX 16, 17	INJECTOR 0.25 MG OR 0.5 MG (2
NINLARO	NEXAVAR 16, 17	`
NIVESTYM INJECTION       98       ML), 2 MG/DOSE (8 MG/3 ML)       61         NOXAFIL ORAL SUSPENSION       121       P         NUBEQA       16, 17       PANRETIN       16, 17         NUCALA       99       pazopanib       16, 17         NUEDEXTA       100       PEMAZYRE       16, 17         NUPLAZID       101       perphenazine-amitriptyline       10         O       phenobarbital       119		
NOXAFIL ORAL SUSPENSION       121       P         NUBEQA       16, 17       PANRETIN       16, 17         NUCALA       99       pazopanib       16, 17         NUEDEXTA       100       PEMAZYRE       16, 17         NUPLAZID       101       perphenazine-amitriptyline       10         O       phenobarbital       119		<i>**</i>
NUBEQA       16, 17       PANRETIN       16, 17         NUCALA       99       pazopanib       16, 17         NUEDEXTA       100       PEMAZYRE       16, 17         NUPLAZID       101       perphenazine-amitriptyline       10         O       phenobarbital       119		
NUCALA		PANRETIN
NUEDEXTA		
NUPLAZID		
O phenobarbital		•
•		
		•

PIQRAY ORAL TABLET 200 MG/DAY	REPATHA SURECLICK
(200 MG X 1), 250 MG/DAY (200 MG	REPATHA SYRINGE
X1-50 MG X1), 300 MG/DAY (150 MG	RETACRIT INJECTION SOLUTION
X 2) 16, 17	10,000 UNIT/ML, 2,000 UNIT/ML,
pirfenidone oral capsule50	20,000 UNIT/ML, 3,000 UNIT/ML,
pirfenidone oral tablet 267 mg, 534 mg, 801	4,000 UNIT/ML, 40,000 UNIT/ML 135
mg	RETEVMO ORAL CAPSULE 40 MG, 80
PLEGRIDY SUBCUTANEOUS PEN	MG
INJECTOR 125 MCG/0.5 ML	REVLIMID
PLEGRIDY SUBCUTANEOUS SYRINGE	REXULTI ORAL TABLET 136
125 MCG/0.5 ML	REZLIDHIA 16, 17
POMALYST	REZUROCK 137
posaconazole oral 121	RINVOQ ORAL TABLET EXTENDED
PRADAXA ORAL PELLETS IN PACKET	RELEASE 24 HR 15 MG, 30 MG, 45
	MG
magabalin aral agrayla 100 mg, 150 mg	RISPERDAL CONSTA
pregabalin oral capsule 100 mg, 150 mg,	
200 mg, 225 mg, 25 mg, 300 mg, 50 mg,	risperidone microspheres
75 mg	roflumilast
pregabalin oral solution	ROZLYTREK ORAL CAPSULE 100 MG,
PREMARIN ORAL	200 MG
PREMPRO	ROZLYTREK ORAL PELLETS IN
PREVYMIS ORAL 125	PACKET
PROCRIT INJECTION SOLUTION 10,000	RUBRACA
UNIT/ML, 2,000 UNIT/ML, 20,000	rufinamide oral suspension
UNIT/ML, 3,000 UNIT/ML, 4,000	rufinamide oral tablet
UNIT/ML, 40,000 UNIT/ML 126	RYDAPT 16, 17
PROLASTIN-C INTRAVENOUS	S
SOLUTION 127	sapropterin141
PROLIA 128	SCEMBLIX ORAL TABLET 20 MG, 40
PROMACTA 129	MG 16, 17
promethazine oral	SECUADO
promethazine rectal suppository 12.5 mg, 25	SIGNIFOR
mg 130	sildenafil (pulm.hypertension) oral tablet 117
promethegan rectal suppository 25 mg, 50	SIRTURO ORAL TABLET 100 MG 144
mg 130	SKYRIZI SUBCUTANEOUS PEN
protriptyline131	INJECTOR
PURIXAN	SKYRIZI SUBCUTANEOUS SYRINGE
pyrimethamine	150 MG/ML 145
Q	SKYRIZI SUBCUTANEOUS WEARABLE
QINLOCK 16, 17	INJECTOR145
quinine sulfate	sodium oxybate
R	sofosbuvir-velpatasvir
REBIF (WITH ALBUMIN) 73	SOMAVERT147
REBIF REBIDOSE73	sorafenib
REBIF TITRATION PACK73	SPRITAM
REPATHA PUSHTRONEX134	

SPRYCEL ORAL TABLET 100 MG, 140	THALOMID 16, 17
MG, 20 MG, 50 MG, 70 MG, 80 MG . 16,	thioridazine 65
17	TIBSOVO 16, 17
STELARA SUBCUTANEOUS 149	tigecycline162
STIVARGA 16, 17	TOBI PODHALER 163
sunitinib malate	tobramycin in 0.225 % nacl 164
SUTENT 16, 17	tolcapone165
SYMDEKO	TRELSTAR INTRAMUSCULAR
SYMPAZAN151	SUSPENSION FOR
SYNAREL 152	RECONSTITUTION 166
SYNRIBO16, 17	tridacaine87
T	trientine oral capsule 250 mg 167
TABLOID 16, 17	trihexyphenidyl
TABRECTA 16, 17	trimipramine
tacrolimus topical153	TRINTELLIX 170
tadalafil (pulm. hypertension) 117	TRULICITY 61
TADLIQ117	TRUQAP16, 17
TAFINLAR 16, 17	TRUSELTIQ ORAL CAPSULE 100
TAGRISSO16, 17	MG/DAY (100 MG X 1), 125
TAKHZYRO154	MG/DAY(100 MG X1-25MG X1), 50
TALTZ AUTOINJECTOR155	MG/DAY (25 MG X 2), 75 MG/DAY (25
TALTZ SYRINGE155	MG X 3)16, 17
TALZENNA ORAL CAPSULE 0.1 MG,	TUKYSA ORAL TABLET 150 MG, 50
0.25 MG, 0.35 MG, 0.5 MG, 0.75 MG, 1	MG16, 17
MG 16, 17	TURALIO ORAL CAPSULE 125 MG, 200
TARGRETIN TOPICAL 156	MG16, 17
TASIGNA	TYMLOS116
tazarotene topical cream 157	$\mathbf{U}$
tazarotene topical gel157	UBRELVY171
TAZORAC TOPICAL CREAM 0.05 % 157	UDENYCA118
TAZORAC TOPICAL GEL 157	UDENYCA AUTOINJECTOR 118
TAZVERIK 16, 17	UDENYCA ONBODY118
TEPMETKO 16, 17	UZEDY139
teriflunomide	V
teriparatide subcutaneous pen injector 20	VALCHLOR172
mcg/dose (620mcg/2.48ml) 116	vancomycin oral capsule 125 mg, 250 mg
testosterone cypionate intramuscular oil 100	
mg/ml, 200 mg/ml159	VANFLYTA16, 17
testosterone enanthate 159	VEMLIDY 175
testosterone transdermal gel in metered-dose	VENCLEXTA 16, 17
pump 159, 160	VENCLEXTA STARTING PACK 16, 17
testosterone transdermal gel in packet 159,	VERQUVO176
160	VERZENIO 16, 17
testosterone transdermal solution in metered	VIBERZI177
pump w/app 159	
tetrabenazine	

VICTOZA SUBCUTANEOUS PEN	XIFAXAN183
INJECTOR 0.6 MG/0.1 ML (18 MG/3	XOLAIR 184
ML)61	XOSPATA 16, 17
VIIBRYD ORAL TABLET 178	XPOVIO ORAL TABLET 100 MG/WEEK
VIIBRYD ORAL TABLETS,DOSE PACK	(50 MG X 2), 40 MG/WEEK (40 MG X
10 MG (7)- 20 MG (23)178	1), 40MG TWICE WEEK (40 MG X 2),
vilazodone	60 MG/WEEK (60 MG X 1), 60MG
VIMPAT ORAL TABLET14	TWICE WEEK (120 MG/WEEK), 80
VITRAKVI ORAL CAPSULE 100 MG, 25	MG/WEEK (40 MG X 2), 80MG TWICE
MG 16, 17	WEEK (160 MG/WEEK)17
VITRAKVI ORAL SOLUTION 16, 17	XTANDI
VIZIMPRO 16, 17	XYREM185
VONJO 16, 17	$\mathbf{Y}$
voriconazole intravenous 179	YONSA17
VOTRIENT 16, 17	${f Z}$
VRAYLAR ORAL CAPSULE 107	ZARXIO186
VRAYLAR ORAL CAPSULE,DOSE	ZEJULA ORAL CAPSULE 17
PACK107	ZEJULA ORAL TABLET 100 MG, 200
$\mathbf{W}$	MG, 300 MG 17
WELIREG 16, 17	ZELBORAF17
X	ZEPATIER187
XALKORI ORAL CAPSULE 16, 17	ZIEXTENZO118
XALKORI ORAL PELLET 150 MG, 20	ziprasidone mesylate57
MG, 50 MG 16, 17	ZOLINZA 17
XATMEP180	ZYDELIG 17
XELJANZ ORAL SOLUTION 181	ZYKADIA 17
XELJANZ ORAL TABLET 181	ZYPREXA RELPREVV
XELJANZ XR 181	INTRAMUSCULAR SUSPENSION
XERMELO 16, 17	FOR RECONSTITUTION 210 MG 188
XGEVA 182	