

Medi-Cal HEDIS® Performance Summary

Medi-Carriebio® renormance duminary					
HEDIS ® Measures for Medi-Cal	HEDIS MY 2021	HEDIS MY 2022	HEDIS MY 2023	HEDIS MY 2023 Percentile (Plan Rating)	Percentile movement
Adherence to Antipsychotic Medications for Individuals with Schizophrenia	64.34%	72.61%	72.69%	90 th (5)	\leftrightarrow
Advising Smokers and Tobacco Users to Quit (CAHPS)	NA Not reported due to small sample size				
Antidepressant Medication Management Effective Continuation Phase Treatment	69.05%	63.39%	56.74%	66.67 th (4)	\downarrow
Appropriate Testing for Pharyngitis	19.49%	18.34%	23.50%	< 10 th (1)	\leftrightarrow
Appropriate Treatment for Upper Respiratory Infection	85.23%	85.23%	82.65%	10 th (2)	\leftrightarrow
Asthma Medication Ratio	59.08%	65.87%	64.98%	33.33rd (3)	\leftrightarrow
Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis ~	40.15%	42.26%	41.93%	< 10 th (1)	\leftrightarrow
Blood Pressure Control for Patients With Diabetes	59.61%	66.42%	70.02%	66.67 th (4)	↑
Breast Cancer Screening	57.84%	58.73%	62.39%	66.67 th (4)	\leftrightarrow
Cervical Cancer Screening	54.01%	56.97%	65.93%	66.67 th (4)	1
Child & Adolescent Well-Care Visits	47.90%	46.78%	51.49%	33.33rd (3)	\leftrightarrow
Childhood Immunization Status Combination 10	28.71%	28.95%	22.99%	10 th (2)	↓
Chlamydia Screening in Women	64.97%	64.88%	67.93%	90 th (5)	1
Controlling High Blood Pressure	60.83%	65.32%	67.55%	66.67 th (4)	1
Diabetes Screening for People with Schizophrenia or Bipolar Disorder who are Using Antipsychotic Medication	83.64%	76.27%	78.21%	33.33 rd (3)	1
Eye Exam for Patients With Diabetes	58.39%	55.23%	61.92%	66.67 th (4)	1
Follow-Up Care for Children Prescribed ADHD Medication Continuation & Maintenance Phase	53.09%	55.20%	50.11%	10 th (2)	↓
Follow-Up after Emergency Department Visit for Mental Illness					
7-day Follow-Up	41.56%	41.72%	48.37%	66.67 th (4)	↑
30-day Follow-Up	57.62%	59.48%	65.71%	66.67 th (4)	↑
Follow-Up After Emergency Department Visit for Substance Use			,		
7-day Follow-Up	3.10%	20.92%	25.05%	Reported as "N	
30-day Follow-Up	6.20%	38.27%	37.53%	NCQ	A
Hemoglobin A1c Control for Patients With Diabetes HbA1c Control < 8%	49.88%	56.20%	57.25%	66.67 th (4)	\leftrightarrow
Immunizations for Adolescents Combination 2	30.41%	34.55%	37.96%	33.33 rd (3)	\leftrightarrow
Kidney Health Evaluation for Patients with Diabetes (KED)	42.95%	45.47%	48.27%	90 th (5)	↑
Lead Screening in Children	46.72%	52.07%	52.39%	10 th (2)	\leftrightarrow



Medi-Cal HEDIS® Performance Summary

medi dai riebio i criormanoc daminary					
HEDIS ® Measures for Medi-Cal	HEDIS MY 2021	HEDIS MY 2022	HEDIS MY 2023	HEDIS MY 2023 Percentile (Plan Rating)	Percentile movement
Metabolic Monitoring for Children and				·	
Adolescents on Antipsychotics	39.29%	36.03%	37.12%	33.33 rd (3)	\leftrightarrow
Blood Glucose and Cholesterol Testing	00.2070	00.0070	0111270	(0)	
Pharmacotherapy for Opioid Use Disorder	23.11%	17.44%	20.30%	10 th (2)	\leftrightarrow
Pharmacotherapy Management of COPD	20.1170	17.4470	20.0070	10 (2)	. ,
Exacerbation					
Systemic Corticosteroid	74.02%	77.35%	79.58%	66.67 th (4)	\leftrightarrow
Bronchodilator	90.59%	89.43%	90.99%	90 th (5)	↑
Plan All-Cause Readmissions Observed-to-Expected Ratio^ - 18-64 Years	0.8802	0.8783	0.8969	66.67 th (4)	↓
Prenatal and Postpartum Care			l		
Timeliness of Prenatal Care	83.94%	88.15%	86.74%	33.33 rd (3)	\downarrow
Timely Postpartum Care	80.05%	79.63%	81.72%	66.67 th (4)	↑
Prenatal Immunization Status Combination	17.38%	14.48%	16.90%	33.33 rd (3)	1
Risk of Continued Opioid Use^ ≥31 days covered	8.59%	6.76%	5.31%	Not reported to NCQA	
Statin Therapy for Patients with Cardiovascular Disease					
Received Statin Therapy	78.77%	80.74%	82.59%	66.67 th (4)	↑
Adherence 80%	85.43%	80.70%	78.58%	66.67 th (4)	\leftrightarrow
Statin Therapy for Patients with Diabetes					
Received Statin Therapy	68.82%	70.11%	70.95%	66.67 th (4)	\leftrightarrow
Adherence 80%	84.05%	76.84%	75.92%	66.67 th (4)	\leftrightarrow
Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics	44.80%	47.96%	47.94%	10 th (2)	\leftrightarrow
Use of Imaging Studies for Low Back Pain~	76.82%	75.90%	73.12%	33.33 rd (3)	↓
Use of Opioids at High Dosage^	2.16%	4.99%	4.17%	Not reported	to NCQA
Use of Opioids from Multiple Providers^ Multiple Prescribers and Multiple Pharmacies	3.41%	3.54%	4.25%	Not reported to NCQA	
Weight Assessment, Counseling for Nutrition and Physical Activity for Children BMI Total	84.43%	82.22%	89.61%	66.67 th (4)	↑
Well-Child Visits in the First 30 Months of Life					
First 15 Months	29.52%	55.79%	59.95%	33.33 rd (3)	\leftrightarrow
15 Months–30 Months	54.93%	62.93%	67.15%	33.33 rd (3)	↑

All benchmarks noted in this report are based on the 2023 NCQA Health Plan Ratings Percentiles and the 2023 NCQA Quality Compass National Benchmarks for Medicaid

~ Inverted rate

Performance Summary Keys:

^{*} Rate calculated per 1000 Members ^ Lower rate is better



Medi-Cal HEDIS® Performance Summary

	Indicates a decrease in the percentile ranking as
↓	compared to the previous measurement year. Each
	arrow down reflects a level of change in the
	percentile ranking. (e.g. a measure that went from
	the 75th percentile to the 50th percentile will
	display one down arrow)
↑	Indicates an increase in the percentile ranking as
	compared to the previous measurement year. Each
	arrow up reflects a level of change in the percentile
	ranking. (e.g. a measure that went from the 25th
	percentile to the 75th percentile will display two
	up arrows)
	Indicates no change in the percentile ranking as
\longleftrightarrow	compared to the previous measurement year.

Plan Rating	National Percentile Range
1	≤10 th percentile
2	>10 th and <33 rd percentile
3	>33 rd and ≤66 th percentile
4	>66 th and ≤90 th percentile
5	>90 th percentile